



SEXING BIRDS USING DISCRIMINANT FUNCTION ANALYSIS: A CRITICAL APPRAISAL

FRANÇOIS-XAVIER DECHAUME-MONCHARMONT,^{1,3} KARINE MONCEAU,¹
AND FRANK CEZILLY^{1,2}

¹Université de Bourgogne, Equipe Ecologie Evolutive, UMR CNRS 5561 Biogéosciences, Dijon, France; and

²Institut Universitaire de France

ABSTRACT.—Discriminant function analysis (DFA) based on morphological measurements is a quick, inexpensive, and efficient method for sex determination in field studies on cryptically monomorphic bird species. However, behind the apparent standardization and relative simplicity of DFA lie subtle differences and pitfalls that have been neglected in some studies. Most of these concerns directly affect assessment of the discriminant performance, a parameter of crucial importance in practice because it provides a measure of the quality of an equation that may be used in later field studies. Using results from 141 published studies and simulations based on a large data set collected on adult Zenaida Doves (*Zenaida aurita*), we assessed the effects of sexual dimorphism, sample size, and validation methods on discrimination rates. We compared the three most common methods used to estimate the proportion of correctly classified males and females by DFA: resubstitution, jackknife, or sample splitting. Results from simulations indicate that these procedures may lead to opposite conclusions, especially when the sample size is small. In particular, the resubstitution techniques appear to be over-optimistic, and we therefore recommend that DFA accuracy be estimated by the jackknife cross-validation procedure. In addition, we show that most previous studies failed to present DFA accuracy with 95% confidence intervals, which hampers comparisons among studies. Finally, our results suggest that large sample sizes should be preferred over repeated measurements of the same individuals, because random measurement error is likely to have only a weak effect on the accuracy of the discriminant rate. *Received 17 May 2010, accepted 11 November 2010.*

Key words: cross-validation, DFA, measurement errors, morphological measurements, sample size effect, sexual dimorphism, *Zenaida aurita*.

Identification du sexe des oiseaux par analyse factorielle discriminante: une évaluation critique

RÉSUMÉ.—L'analyse factorielle discriminante (AFD) à partir de mesures morphologiques constitue une méthode rapide, peu chère et efficace pour un sexage sur le terrain d'oiseaux appartenant à des espèces cryptiquement monomorphes. Cependant, l'apparente standardisation et la relative simplicité de l'AFD cachent en réalité plusieurs différences et écueils subtils négligés dans de nombreuses études. La plupart de ces problèmes nuisent directement à l'évaluation du pouvoir de discrimination, un paramètre crucial en pratique puisqu'il correspond à une mesure de la qualité de l'équation destinée à être utilisée dans des études de terrain ultérieures. En nous appuyant sur 141 publications et sur des simulations basées sur un large jeu de données collectées sur des Tourterelles à queue carrée *Zenaida aurita* adultes, nous avons évalué l'effet du dimorphisme sexuel, de la taille de l'échantillon et de la méthode de validation sur le taux de discrimination. Nous avons comparé les trois méthodes les plus utilisées pour estimer la proportion de mâles et de femelles correctement sexés par l'AFD: la resubstitution, le jackknife et le sample-splitting. Les résultats issus des simulations indiquent que ces procédures peuvent conduire à des conclusions opposées, surtout en cas de petites tailles d'échantillons. Plus précisément, les techniques de resubstitution semblent beaucoup trop optimistes. Nous recommandons que la précision d'une AFD soit évaluée au moyen d'un jackknife. De plus, nous montrons que la plupart des études antérieures indiquent cette précision sans l'assortir d'un intervalle de confiance à 95%, ce qui limite la possibilité de comparaisons entre études. Enfin, nos résultats suggèrent que de grandes tailles d'échantillons devraient être préférées à des mesures répétées sur les mêmes individus puisqu'il est probable que l'erreur de mesure n'ait qu'un effet limité sur la précision du taux de discrimination.

³E-mail: fx.dechaume@u-bourgogne.fr

SEX-RELATED DIFFERENCES in the behavior and ecology of bird species are crucial to our understanding of sexual selection and mating systems (Andersson 1994) and may also have important consequences for management and conservation (Zavalaga and Paredes 1997, Fernandez-Juricic et al. 2009). Therefore, the ability to identify the sex of an individual is of paramount importance in avian studies. Although sexual dimorphism can be conspicuous (e.g., plumage-based sexual dichromatism), especially in the case of polygynous species (Darwin 1871, Andersson 1994), differences between male and female birds are often subtle or escape the human eye (Cuthill et al. 1999). Several techniques have been developed to alleviate this difficulty, including anatomical examination (Petrides 1950, Miller and Wagner 1955), vocalization analyses (Bourgeois et al. 2007), sex-specific behavior analyses (Castoro and Guhl 1958, Flux and Innes 2001, Fletcher and Hamer 2003), and, more recently, molecular techniques (Dubiec and Zagalska-Neubauer 2006). Although the latter methods are by far the most popular techniques and are thought to be very reliable (but see Robertson and Gemmell 2006), they require training and a license to collect blood or tissues, which raises ethical issues, and imply financial costs and delays for processing of samples.

Sexing based on morphometrics is often a reasonable choice for quick and inexpensive but efficient sex identification in field studies on cryptically monomorphic bird species. Significant, albeit small, differences in biometric measurements often exist between females and males, offering the possibility of discriminating between the sexes (Murphy 2007, Cardoni et al. 2009). To that end, various statistical analyses have been used, such as linear models (Jeffrey et al. 1993, Iko et al. 2004, Ura et al. 2005, Gill and Vohnhof 2006, Hallgrimsson et al. 2008) or multivariate methods, including principal component analysis (Rubega 1996, McCracken et al. 2000, Remisiewicz and Wennerberg 2006, Urfi and Kalam 2006, Schroeder et al. 2008) and discriminant function analysis (DFA). The latter is the most popular of these statistical methods, and its use has increased steadily in recent decades (Fig. 1). The principle

of DFA is to provide equations based on morphological measurements in order to predict the sex of birds. This equation is primarily calibrated on individuals of known sex. Every discriminant equation comes with its estimated proportion of correctly sexed individuals.

Biologists justifiably pay great attention to the proportion of misclassified birds, because a discriminant equation is not published as an end in itself but as a tool for reliably and rapidly sexing birds in later field studies. Various methods can be used to estimate the proportion of correctly classified males and females by DFA, and yet the choice of a method is rarely justified in the avian literature. In addition, the estimated rate of correct discrimination has been reported to be sensitive to sample size (Morrison 1984). Among or within subjects, variance in measurements may also have an effect on the ability to discriminate males from females. Such variance has two components: the true variance among individuals and the measurement error (ME) that arises from random or systematic errors. Large MEs increase the risk of Type II error (Lougheed et al. 1991, Yezzerinac et al. 1992). Francis and Matlin (1986) showed that discriminant power could fall from 89% to <50% when a small amount of bias exists in morphometrics measurement. The need for ME assessment has been emphasized (Bailey and Byrnes 1990, Arnqvist and Mårtensson 1998), yet few studies that used DFA have reported ME (Flux and Innes 2001, Devlin et al. 2004, Kenward et al. 2004), and its potential effect on discrimination rate has rarely been investigated (but see Mallory and Forbes 2005).

Here, we address the problem using a twofold approach. First, using simulations based on a large data set, we assess the choice of validation method and the effect of sample size or measurement errors on the estimated accuracy of the DFA. Second, we review the literature to quantify the methodologies used, assess the effects of the extent of sexual dimorphism, sample size, or number of variables on discriminant rate, and test predictions from our simulations. Finally, we provide recommendations for future studies.

METHODS

Simulations.—A total of 525 adult Zenaida Doves (*Zenaida aurita*; 294 females and 231 males) were captured on Barbados from March to May 2007 (Monceau et al. 2011). This sexually monomorphic species is widely distributed throughout the Caribbean. Each individual was molecularly sexed (Fridolfsson and Ellegren 1999). The reliability of molecular sexing techniques has been reported to vary according to which PCR amplification procedure is used (Dubiec and Zagalska-Neubauer 2006, Robertson and Gemmell 2006, Daniel et al. 2007). In a preliminary study (Monceau 2009), we first assessed the reliability of our procedure by comparing results from molecular sexing with behavioral observations based on 48 pairs.

For each bird, we measured three dimensions of the bill at nostrils (length, depth, and width), head plus bill length, left and right tarsus length, left and right wing chord, and tail length. All measurements were made by the same person (K.M.) with a digital caliper (precision: ± 0.2 mm), except for wing chord and tail length, which were measured with a ruler (precision: ± 1 mm). Each character was measured twice, and the caliper was removed between measurements in order to assess MEs (Bailey and Byrnes 1990).




FIG. 1. Histogram of the number of papers in our survey that reported sexing birds by means of discriminant function analysis ($n = 141$ articles) over the past 50 years.

TABLE 1. Comparisons between male and female Zenaida Doves on Barbados for all characters with measurement error (ME) and mean difference index (MDI) calculated as (mean female/mean male) \times 100.

Characters	Males (mean \pm SD)	Females (mean \pm SD)	Cohen's d (95% CI)	<i>t</i> -test	<i>P</i>	ME (%)	MDI (%)
Bill length (mm)	10.67 \pm 0.48	10.45 \pm 0.51	0.45 (0.28–0.62)	–5.11	$<10^{-5}$	7.92	97.94
Bill width (mm)	4.07 \pm 0.22	3.93 \pm 0.26	0.58 (0.41–0.77)	–6.73	$<10^{-5}$	9.63	96.56
Bill depth (mm)	4.15 \pm 0.21	3.98 \pm 0.21	0.78 (0.61–0.96)	–8.83	$<10^{-5}$	21.86	95.90
Head plus bill length (mm)	48.59 \pm 0.99	47.40 \pm 1.02	1.18 (1.01–1.36)	–13.47	$<10^{-5}$	3.65	97.55
Mean tarsus length (mm)	26.76 \pm 0.86	25.94 \pm 0.77	1.01 (0.92–1.21)	–11.30	$<10^{-5}$	6.83	96.94
Mean wing chord (cm)	15.31 \pm 0.54	14.74 \pm 0.46	1.61 (0.98–1.36)	–12.93	$<10^{-5}$	2.22	96.28
Tail length (cm)	10.38 \pm 0.56	9.94 \pm 0.56	0.79 (0.60–0.97)	–8.94	$<10^{-5}$	4.07	95.76
Body mass (g)	149.19 \pm 15.21	141.81 \pm 13.95	0.51 (0.33–0.70)	–5.73	$<10^{-5}$	—	95.05

Because MEs were low for all traits except bill depth (Table 1), we used the mean of first and second measurements of each variable in subsequent analyses. Birds were weighed with a Pesola digital pocket scale MS500 (precision: ± 0.1 g). Males differed significantly from females in all morphological traits, and the mean difference index (MDI = $100 \times$ mean female/mean male; Delestrade 2001, Helfenstein et al. 2004) was $>95\%$ for all characters (Table 1).

We performed DFA on the original data set using the eight morphological variables. Because of a violation of the assumption of homogeneity of the variance–covariance matrices (box's *M* test, $\chi^2 = 267.1$, *df* = 190, $P = 0.0002$), we performed quadratic DFA (Stevens 1992) using the *qda* function from the MASS package (Venables and Ripley 2002) for R, version 2.10.1 (R Development Core Team 2010). We searched for potential multivariate outliers by measuring robust Mahalanobis distances (Rasmussen 1988, Jarrell 1994, Tabachnick and Fidell 2000, Nordhausen et al. 2008). We assessed the leverage of outliers by removing them from the data set and estimating the new discriminant rate. The effect of automated variable selection was assessed using the *stepclass* function and minimization of Wilks's lambda criterion (Mardia et al. 1979) as implemented in the *klaR* package for R (Weihs et al. 2005).

We compared three validation methods commonly used in sexing birds to estimate the proportion of correctly classified individuals by a DFA: resubstitution, jackknife (Manly 1994), and sample splitting (Picard and Berk 1990). In the first method, the sex of each individual is predicted using the discriminant function calculated from the complete data set. Using the jackknife (or leave-one-out) method, the sex of an individual is predicted from the discriminant equation calculated after that individual has been removed from the data set. This procedure is repeated until a sex is assigned to each individual (Tabachnick and Fidell 2000). With sample splitting, the data set is randomly divided into two subsamples, and the training set (two-thirds of the individuals, as recommended by Picard and Berk 1990) is used to compute the discriminant function that is then used to assess the accuracy of the discriminant function by classifying the remaining third of the individuals.

We simulated both smaller data sets and data sets with larger MEs in order to assess the effect of sample size, the choice of validation method, and MEs on the discriminant power. We defined 100 different sample sizes ranging from 25 to 520 individuals, regularly spaced every five individuals. For each sample size, we simulated 500 different data sets by randomly sampling individuals

from the complete data set. For each of these 50,000 simulated data sets, we performed DFA and assessed the discriminant rates using resubstitution, jackknife, and sample-splitting methods.

We performed a second series of simulations using the complete original data set ($n = 525$ individuals) to assess the effect of ME. Each variable used in the DFA was the mean of two repeated measures recorded in order to assess ME (Table 1). We simulated higher ME for a given variable by adding random noise to both repeated measures. This additional noise followed a normal distribution with mean $\mu = 0$ and standard deviation σ . We adjusted σ up to obtain the chosen ME ($\pm 0.01\%$). Note that it was not possible to artificially decrease ME, because we could not improve the accuracy of the data. Variables with ME already higher than the ones requested for the simulation were not modified. For example, for a minimal ME of 10%, we increased ME for every variable (including mass), except for bill depth, which already showed a larger ME (Table 1). For each value of minimal ME, we simulated 500 data sets and performed a DFA on each data set. For each DFA, the proportion of correctly sexed individuals was estimated using the jackknife procedure.

Literature survey.—We searched the literature and found 141 studies (from 1961 to 2010) on 132 species from 43 families (online supplement; see Acknowledgments) that used DFA to identify an individual's sex or describe sexual dimorphism in birds. For a given group (same population and same age class), some authors reported more than one discriminant rate using several discriminant equations or several validation methods or both. In order to avoid giving too much weight to studies that reported multiple DFAs, we retained only the equation that led to the best discrimination between sexes. If the resubstitution method or sample-splitting was used along with the jackknife, we report only the discriminant rate estimated for the jackknife validation procedure. We collected a total of 186 DFAs on different populations, age classes, and species. For each of these analyses, we recorded the sample size used to calculate the discriminant function, the validation method, and the discriminant rate. Some authors used a stepwise procedure to reduce the number of variables used in the discriminant equation. We reported the stepwise procedure used (if available) and the number of variables retained in the discriminant equation. Finally, the mean tarsus length of males and females was also collected, if available, in order to estimate the relative sexual dimorphism (Storer's index: absolute value of the difference between male and female divided by the mean tarsus

length; Smith 1999). We assessed the effect of relative sexual dimorphism on the number of birds sampled using analysis of variance and on the number of variables selected in the discriminant equation using generalized linear models (Poisson regression). We used linear models to assess the effect on the discriminant rate of sample size, relative sexual dimorphism, validation method, number of variables in the discriminant equation, variable-selection methods (automated or not), and year of publication. Percentages were inspected for normality and normalized using square-root arcsine transformation if needed (Zar 1999).

RESULTS

Effect of sample size and validation methods.—The differences between the three validation methods were far from negligible, especially for small sample sizes (Fig. 2). The proportions of correctly sexed birds (discriminant rate) were larger when estimated with the resubstitution method, and decreased with increasing sample size (Fig. 2A). For small data sets ($n < 60$ birds), the mean estimate of the discriminant rate was $>90\%$. By contrast, the mean estimates with the jackknife (Fig. 2B) or sample-splitting (Fig. 2C) methods increased with increasing sample size and were consistent, although the variance of the estimates was smaller with the jackknife than with the sample-splitting method.

Differences between methods were less pronounced with large data sets. Considering the DFA based on the complete original data set ($n = 525$ adults), the resubstitution method estimated a proportion of correctly sexed adults of 81.0% (95% confidence interval [CI]: $77.5\text{--}84.8\%$), whereas the same value was 80.4% (95% CI: $76.8\text{--}84.4\%$), and 79.1% (95% CI: $75.6\text{--}82.6\%$) for the jackknife and the sample-splitting methods, respectively. Because their bootstrapped 95% CI largely overlapped, the estimated proportion of correctly classified birds did not significantly differ according to the validation method. By measuring robust Mahalanobis distances, we identified 39 potential multivariate outliers (19 males and 20 females). However, these outliers had a limited effect in our analyses, because removing these extreme individuals from the original data set only slightly increased the proportion of correctly sexed birds estimated with a given validation method: resubstitution, 84.0% (95% CI: $80.0\text{--}87.7\%$); jackknife, 82.0% (95% CI: $77.7\text{--}86.0\%$); and sample splitting, 80.1% (95% CI: $73.7\text{--}85.7\%$).

The stepclass function for automated variable selection suggested several subsets of variables. The most frequent set was made up of three variables: wing chord, head length, and tarsus length. Stepwise variable selection based on minimization of Wilks's lambda criterion, a frequently used procedure in the bird literature, led to much larger variable sets, with up to eight variables being included in the complete sample ($n = 525$). We used the reduced set of three variables obtained from the stepwise procedure (wing chord, head length, and tarsus length) and performed the same simulations detailed above. The shape of the relationship between mean discriminant rate and sample size was unchanged, though it was slightly less steep for small sample sizes. In addition, asymptotic discriminant rates calculated for the complete sample size ($n = 525$) were slightly smaller, with 79.8% (95% CI: $74.8\text{--}82.5\%$), 78.9% (95% CI: $74.2\text{--}81.9\%$), and 78.1% (95% CI: $74.9\text{--}86.1\%$) of correctly sexed birds for resubstitution, jackknife, and sample-splitting, respectively.




FIG. 2. Simulation of the effect of sample size on the estimated proportion of correctly classified individuals (discriminant rate) in simulated discriminant function analyses (DFAs) of the Zenaida Dove data from Barbados. From the complete data set ($n = 525$ individuals), smaller subsamples were randomly selected (ranging from 25 to 520 individuals, with 500 subsamples per size). For each of these 50,000 subsamples, we performed a DFA and evaluated the discriminant rate by three methods: (1) resubstitution; (2) jackknife cross-validation; and (3) sample-splitting, in which two-thirds of the data set was used as the training sample and the remaining third as the test sample. Each gray dot represents one DFA. Each dot was slightly randomly jittered to reduce overplotting. The thick line and the dotted lines, respectively, represent the mean discriminant rate and the 95% limits computed from the 500 DFA performed for each sample size. The horizontal dashed line represents the discriminant rate estimated from the complete data set.

Effect of measurement errors.—An increase of ME led to a decrease in the proportion of correctly sexed individuals (Fig. 3). Despite a large drop of accuracy in the simulated measurements, the mean proportion of correctly classified individuals only decreased from 80.38% to 78.09% , which corresponds to ~ 12 additional adults incorrectly classified. Using the smaller subset of variables (wing chord, head length, and tarsus length) selected by automated stepwise procedure, we found similar results. The mean discriminant rate dropped marginally from 78.9% to 77.7% with a simulated ME of 30% .

Literature survey.—Sample size varied greatly, from 10 birds (4 males and 6 females) to 1,891 (918 males and 973 females). The median sample size was 80 birds (95% CI: $72.5\text{--}94.0$). The median sex ratio (male/female) was 1.04 (95% CI: $0.97\text{--}1.09$). Discriminant




FIG. 3. Boxplot of the jackknifed estimated discriminant rate (proportion of correctly classified individuals) as a function of the minimal measurement error (ME). ME was increased to a given value by adjusting random noise to the repeated measures except when real ME was already larger. For each value of minimal ME, 500 data sets were simulated from the complete data set. Every discriminant equation relied on the eight variables listed in Table 1. For a given minimal ME, the black circle depicts the mean discriminant rate, the thick line the median, and the box the interquartile range. The horizontal dashed line represents the reference discriminant rate estimated from the original data, with no noise added.

rates ranged from 63% to 100% (median = 91.8%, 95% CI: 90.3–93.2). Among the 186 DFAs, 40.9% relied on the resubstitution validation method or simply reported the proportion of birds correctly sexed (presumably using resubstitution method), whereas the jackknife and sample-splitting methods, respectively, accounted for 39.8% and 19.4%. The first occurrence of the jackknife validation method in our survey dates back to the mid-1980s (Brennan et al. 1984), and one might argue that use of the jackknife is less common in our survey because it is a more recent and computer-intensive method than the other two. However, although most of the modern statistical packages (e.g., R, SAS, and SPSS) implement DFAs with jackknife procedures, only 47.4% of 95 DFA studies published since 2000 relied on jackknife procedures. Among the 186 DFAs, 62.4% used the automated stepwise procedure to reduce the number of variables in the discriminant function. There was no difference in number of variables in the discriminant equation between studies that relied on automated stepwise variable selection and the others (Wilcoxon test, $W = 4,327$, $P = 0.43$). Measurement errors were mentioned in <5% of the papers.

Relative sexual dimorphism had the largest effect on the discriminant rate ($F = 55.69$, $df = 1$ and 122 , $P < 10^{-5}$) and explained 30.4% of the variance. Neither the validation method ($F = 0.25$, $df = 2$ and 127 , $P = 0.78$), the year of publication ($F = 2.81$, $df = 1$ and 128 , $P = 0.096$), nor the variable selection procedure (automated or not) ($F = 1.71$, $df = 1$ and 128 , $P = 0.19$) had an effect on the discriminant rate. However, the interaction between sample size and




FIG. 4. Results of a literature survey on the effect of relative sexual dimorphism (absolute difference of tarsus length between males and females divided by mean tarsus length) on the discriminant rate (proportion of correctly sexed birds). The line and the gray area depict the predicted trend and its 95% confidence interval fitted by linear regression ($F = 64.7$, $df = 1$ and 132 , $P < 10^{-5}$) after square-root arcsine transformation for normalization of the percentages.

the number of variables was significant ($F = 9.16$, $df = 1$ and 129 , $P = 0.003$), with the discriminant rate decreasing with an increasing number of variables for small sample sizes. No other interaction had a significant effect on the discriminant rate (all $P > 0.18$). The discriminant rate significantly increased with the relative sexual dimorphism (Fig. 4). One might expect that the authors adjusted their measurement effort (sample size or number of morphological variables) accordingly. However, there was no significant association between relative sexual dimorphism and number of birds sampled ($F = 1.24$, $df = 1$ and 132 , $P = 0.27$) or the number of variables used in the discriminant equation (Poisson regression: $\chi^2 = 0.27$, $df = 1$, $P = 0.16$).

Contrary to our simulations, validation methods had no significant effect on the discriminant rate in published studies, and lower sample sizes had an effect only in interaction with the number of variables. This might be explained by publication bias. We therefore examined the funnel plots of discriminant rate as a function of the log-transformed sample size (Palmer 2000). Because of sampling error, the variance in estimates of the discriminant rate is expected to be higher for studies with smaller sample sizes (Møller and Jennions 2001). For each validation method, there was no significant difference between the variance of discriminant rates of the smallest 50% of sample sizes and the largest 50% (Levene test; resubstitution: $P = 0.34$, jackknife: $P = 0.69$, sample-splitting: $P = 0.45$), which reveals a lack of variance for small sample sizes. Because there was no significant relationship between sexual dimorphism and sample size, the lack of variance for small sample sizes cannot be attributed to studies on strongly dimorphic species (which require fewer individuals to generate a high discriminant rate) and is possibly, then, attributable to a publication bias. Small data sets that led to small discriminant rates were probably rejected by editors or underreported by the authors.

DISCUSSION

Effect of the validation method.—The simulations based on our Zenaida Dove data set indicate that the choice of validation method may have a strong effect on the estimated discriminant rate. This effect was particularly sensitive for small to intermediate sample sizes (<200 individuals). Estimates of the discriminant rate based on subsampling techniques (such as jackknife cross-validation or splitting of the data into training and test sets) should be preferred over estimates based on resubstitution, which appeared to be overly optimistic for small sample sizes. It has been well illustrated, in both the statistical (Johnson and Wichern 1992, Huberty 1994, Manly 1994, Piraux and Palm 2001, Wehberg and Schumacher 2004) and the applied literature (Eisenbeis 1977, Lance et al. 2000), that reclassification of the original individuals used in constructing the discriminant equation leads to a biased estimated discriminant rate. In one of the first papers to use DFA to sex birds, Ryder (1978) explicitly raised the problem of resubstitution and recommended that it not be used, especially in the case of a comparison among studies. But as the technique became more popular within the ornithologist community, this advice was somehow overlooked. The two subsampling techniques are not equivalent, however. Sample splitting led to a mean estimate of the proportion of correctly classified individuals consistent with the mean estimate using the jackknife procedure, but with a much larger variance. Therefore, jackknife cross-validation should always be preferred over the sample-splitting procedure or resubstitution.

Surprisingly, we found no difference in the discriminant power between these three validation methods in our survey. However, most published discriminant rates were quite high (median discriminant rate = 91.8%). Even in the case of minute differences between males and females (<1% difference), the mean discriminant rate reported in published studies was still ~80%. It is possible that many authors (e.g., Clark et al. 1991) restrained themselves from publishing DFAs with low discriminant rates, which has led to a “file-drawer effect” (Scargle 2000, Møller and Jennions 2001) and underestimation of the influence of the validation method.

Interstudy comparisons.—The authors often compared their discriminant rates to those of previous studies, most of the time to claim that the new discriminant function led to better classification, even if it increased the discriminant rate by only a small percentage. Such narrow differences may be meaningless, especially when different validation techniques were used to estimate the discriminant rate. In addition, all the 141 studies that we reviewed failed to report 95% CIs around discriminant rates, thus hampering comparisons among studies (Nakagawa and Cuthill 2007, Garamszegi et al. 2009), even when conducted on the same species. Just as publishing means without indication of variance is not acceptable, one should never report discriminant rates without their confidence intervals. Bootstrap resampling provides a simple and robust method for calculating this range. Another possibility would be to consider Bayesian posterior probabilities instead of just looking at overall error rates (McCarthy 2007, Garamszegi et al. 2009, Hastie et al. 2009). Another kind of comparison is also highly questionable. Some authors compared apparent error (i.e., estimated from resubstitution procedure) and error estimated by sample-splitting in order to detect possible sampling bias (e.g., Hanners and Patton 1985). As long as these two error rates were

not too different, they concluded that there was no sampling bias. This kind of argument is poorly sustained, because important differences between these two estimates were possible in our simulations even in the absence of sampling bias.

Importance of large sample size.—Our simulations with small to medium sample sizes (<200 birds) highlighted the wide variance in the estimated proportion of misclassified birds. High discriminant success can be obtained by chance, raising reasonable doubts about the subsequent use of equations constructed from small data sets. In accordance with previous studies (Brennan et al. 1991, Wehberg and Schumacher 2004, Shealer and Cleary 2007, Isaksson et al. 2008), we recommend caution in dealing with discriminant equations computed from small data sets. In addition, the sample size in interaction with the number of variables in the equation had a significant effect on the discriminant rates from our literature survey. If there are too few individuals in relation to the number of variables, the analysis leads to a poor discriminant rate (Marks and Dunn 1974, Burnham and Anderson 2002). This result is known as the “curse of dimensionality” (Bellman 1957). Several authors have advised carefully adjusting the number of birds measured to the number of morphological variables in order to have a sample size $\geq 3 \times$ larger than the number of variables used in the DFA (Williams and Titus 1988, McGarigal et al. 2000), but this criterion is sometimes claimed to be arbitrary (James and McCulloch 1990).

Variable selection.—Estimating the optimal number of variables to measure is of obvious importance. Morphological measurements are usually highly correlated, which can lead to unstable parameter estimates. Eliminating multicollinearity among the variables could improve the discriminant rate, although the procedure is highly debated (James and McCulloch 1990). A common procedure is to reduce each subset of highly correlated variables to only one variable. More importantly, morphometric measures are costly in time and can be stressful for the birds. Field biologists have to limit themselves to a narrow set of morphological variables. Two-thirds of the authors relied on automated stepwise techniques in order to identify, from a larger number of potentially relevant measures, a reasonable set of variables. Biologists should be extremely cautious with the output of automated analysis. Several authors strongly recommend that stepwise procedure be avoided entirely (James and McCulloch 1990, Snyder et al. 1991, Thompson 2001). One peer-reviewed journal even made it an editorial policy to summarily reject any article that used the technique (Babyak 2004). Stepwise methods capitalize on sampling error and yield results that are not replicable (Thompson 1995); there is no reason to think that they lead to the best equation for sexing the model species, and no guarantee that the results will be useful for later studies (James and McCulloch 1990, Derksen and Keselman 1992). Most importantly, the greatest danger of the use of stepwise analysis is the temptation to leap directly from routine procedure to straightforward conclusions about ecological relevance. These procedures are unable to select from a set of variables those that are most influential (Burnham and Anderson 2002); therefore, the selected variables are not necessarily more biologically relevant than nonselected ones.

We offer the reminder to authors that stepwise procedures are exploratory and only suggest combinations of variables that do an adequate job, not necessarily the most discriminating or the simplest. Our survey of the literature showed that studies that relied

on automated stepwise procedures did not differ from the others in the number of variables used in the discriminant equation or on their discriminant power. Scientific judgment and choice of biologically meaningful combination of variables must play a leading role in selecting variable (James and McCulloch 1990, Burnham and Anderson 2002). If stepwise procedures are used, Manly (1994) advised that the analyses be rerun several times with randomized subsamples of individuals to check the validity of the results.

Measurement errors.—We also found that the effect of ME on discriminant rates is generally moderate but nonetheless significant. Simulating large ME in our measures only marginally decreased the estimated proportion of correct classification. This result contrasts sharply with that obtained by Francis and Mattlin (1986), who reported high sensitivity of DFA to small amounts of bias in measurement. They simulated systematic bias by adding a constant to each of the dimension of their data set, which was later analyzed with a discriminant function constructed from the unmanipulated data set. We, on the other hand, investigated the effect of random bias. As long as these MEs equally affected males and females in the same proportion and with no systematic bias, they had a limited effect on the probability of misclassification. This suggests that increasing the number of sampled birds might be preferable to additional measurements of the same bird. Yet, because measurement errors were rarely reported in the papers we reviewed, we could not compare our results from the simulation with actual field data. More studies are needed before firm conclusions can be made regarding the importance of ME for the accuracy of DFA.

Finally, several authors warned that multivariate outliers can lead to violation of multinormality assumption and can severely impair the accuracy of DFA (Manly 1994, Tabachnick and Fidell 2000, Osborne and Overbay 2004). Rather extreme observations can inflate classification error rates and bias parameter estimation. However, search methods for outliers were almost never reported in the ornithological literature that we reviewed (but see Blakesley et al. 1990, Santiago-Alarcon and Parker 2007). We advise that future studies that use DFA to sex birds systematically test for the presence of outliers and, if they are present, decide, on the basis of established criteria (discussed at length in Tabachnick and Fidell 2000), whether or not to remove them.

Our purpose was not to criticize the use of DFA, which remains in our opinion a very efficient and straightforward technique. Molecular methods are not foolproof, but they are unquestionably more reliable and popular than DFA. If a person does not have access to a molecular laboratory, proper performance of DFA should follow the guidelines that we have recommended. Most of the common pitfalls addressed here can easily be avoided. Authors should be extremely cautious with the predictions of discriminant equations constructed from small data sets, even when the reported discriminant rate is high. The accuracy of DFA should be estimated with the jackknife cross-validation procedure, and this accuracy should always be presented with 95% CIs to allow interstudy comparisons.

ACKNOWLEDGMENTS

Supplementary material (Table S1) is available at [dx.doi.org/10.1525/auk.2011.10129](https://doi.org/10.1525/auk.2011.10129). We thank S. Devonish of the Natural Heritage Department for providing us with a permit to capture and ring Zenaida Doves in Barbados. We thank J. Moreau, S. Motreuil, and

G. Prato for help in catching birds and W. Worrell for hospitality. K.M. was supported by a doctoral grant from the Ministère de la Recherche et de l'Enseignement Supérieur. Financial support was partly provided by a research grant from the Agence Nationale pour la Recherche (program Monogamix ANR-08-BLAN-0214-02). Calculations were performed using HPC resources from DSI-CCUB (Université de Bourgogne).

LITERATURE CITED

- ANDERSSON, M. 1994. Sexual Selection. Princeton University Press, Princeton, New Jersey.
- ARNQVIST, G., AND T. MÅRTENSSON. 1998. Measurement error in geometric morphometrics: Empirical strategies to assess and reduce its impact on measures of shape. *Acta Zoologica Academiae Scientiarum Hungaricae* 44:73–96.
- BABYAK, M. A. 2004. What you see may not be what you get: A brief, nontechnical introduction to overfitting in regression-type models. *Psychosomatic Medicine* 66:411–421.
- BAILEY, R. C., AND J. BYRNES. 1990. A new, old method for assessing measurement error in both univariate and multivariate morphometric studies. *Systematic Zoology* 39:124–130.
- BELLMAN, R. 1957. Dynamic Programming. Princeton University Press, Princeton, New Jersey.
- BLAKESLEY, J. A., A. B. FRANKLIN, AND R. J. GUTIÉRREZ. 1990. Sexual dimorphism in Northern Spotted Owls from northwest California. *Journal of Field Ornithology* 61:320–327.
- BOURGEOIS, K., C. CURÉ, J. LEGRAND, E. GÓMEZ-DÍAZ, E. VIDAL, T. AUBIN, AND N. MATHEVON. 2007. Morphological versus acoustic analysis: What is the most efficient method for sexing Yelkouan Shearwaters *Puffinus yelkouan*? *Journal of Ornithology* 148:261–269.
- BRENNAN, L. A., J. B. BUCHANAN, C. T. SCHICK, AND S. G. HERMAN. 1991. Estimating sex-ratios with discriminant function analysis: The influence of probability cutpoints and sample size. *Journal of Field Ornithology* 62:357–366.
- BRENNAN, L. A., J. B. BUCHANAN, C. T. SCHICK, S. G. HERMAN, AND T. M. JOHNSON. 1984. Sex determination of Dunlins in winter plumage. *Journal of Field Ornithology* 55:343–348.
- BURNHAM, K. P., AND D. R. ANDERSON. 2002. Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach, 2nd ed. Springer-Verlag, New York.
- CARDONI, D. A., J. E. MALDONADO, J. P. ISACCH, AND R. GREENBERG. 2009. Subtle sexual dimorphism in the Bay-capped Wren-Spintail (*Spartonoica maluroides*; Furnariidae) uncovered through molecular sex determination. *Ornitología Neotropical* 20:347–355.
- CASTORO, P. L., AND A. M. GUHL. 1958. Pairing behavior of pigeons related to aggressiveness and territory. *Wilson Bulletin* 70:57–69.
- CLARK, R. G., P. C. JAMES, AND J. B. MORARI. 1991. Sexing adult and yearling American Crows by external measurements and discriminant analysis. *Journal of Field Ornithology* 62:132–138.
- CUTHILL, I. C., A. T. D. BENNETT, J. C. PARTRIDGE, AND E. J. MAIER. 1999. Plumage reflectance and the objective assessment of avian sexual dichromatism. *American Naturalist* 153:183–200.
- DANIEL, C., C. D. MILLAR, S. M. H. ISMAR, B. M. STEPHENSON, AND M. E. HAUBER. 2007. Evaluating molecular and behavioural sexing methods for the Australasian Gannet (*Morus serrator*). *Australian Journal of Zoology* 55:377–382.

- DARWIN, C. R. 1871. *The Descent of Man, and Selection in Relation to Sex*. John Murray, London.
- DELESTRADE, A. 2001. Sexual size dimorphism and positive assortative mating in Alpine Choughs (*Pyrrhocorax graculus*). *Auk* 118:553–556.
- DERKSEN, S., AND H. J. KESELMAN. 1992. Backward, forward and stepwise automated subset selection algorithms: Frequency of obtaining authentic and noise variables. *British Journal of Mathematical and Statistical Psychology* 45:265–282.
- DEVLIN, C. M., A. W. DIAMOND, AND G. W. SAUNDERS. 2004. Sexing Arctic Terns in the field and laboratory. *Waterbirds* 27:314–320.
- DUBIEC, A., AND M. ZAGALSKA-NEUBAUER. 2006. Molecular techniques for sex identification in birds. *Biological Letters* 43:3–12.
- EISENBEIS, R. A. 1977. Pitfalls in the application of discriminant analysis in business, finance, and economics. *Journal of Finance* 32:875–900.
- FERNANDEZ-JURICIC, E., A. J. DEL NEVO, AND R. POSTON. 2009. Identification of individual and population-level variation in vocalizations of the endangered Southwestern Willow Flycatcher (*Empidonax traillii eximius*). *Auk* 126:89–99.
- FLETCHER, K. L., AND K. C. HAMER. 2003. Sexing terns using biometrics: The advantage of within-pair comparisons. *Bird Study* 50:78–83.
- FLUX, I., AND J. INNES. 2001. A field technique for determining the sex of North Island Kokako (*Callaeas cinerea wilsoni*). *Notornis* 48:217–223.
- FRANCIS, R. I. C. C., AND R. H. MATTLIN. 1986. A possible pitfall in the morphometric application of discriminant analysis: Measurement bias. *Marine Biology* 93:311–313.
- FRIDOLFSSON, A. K., AND H. ELLEGREN. 1999. A simple and universal method for molecular sexing of non-ratite birds. *Journal of Avian Biology* 30:116–121.
- GARAMSZEGI, L. Z., S. CALHIM, N. DOCHTERMANN, G. HEGYI, P. L. HURD, C. JØRGENSEN, N. KUTSUKAKE, M. J. LAJEUNESSE, K. A. POLLARD, H. SCHIELZETH, and others. 2009. Changing philosophies and tools for statistical inferences in behavioral ecology. *Behavioral Ecology* 20:1363–1375.
- GILL, S. A., AND M. J. VONHOF. 2006. Sexing monochromatic birds in the field: Cryptic sexual size dimorphism in Buff-breasted Wrens (*Thryotorus leucotis*). *Ornitología Neotropical* 17:409–418.
- HALLGRIMSSON, G. T., S. PALSSON, AND R. W. SUMMERS. 2008. Bill length: A reliable method for sexing Purple Sandpipers. *Journal of Field Ornithology* 79:87–92.
- HANNERS, L. A., AND S. R. PATTON. 1985. Sexing Laughing Gulls using external measurements and discriminant analysis. *Journal of Field Ornithology* 56:158–164.
- HASTIE, T., R. TIBSHIRANI, AND J. FRIEDMAN. 2009. *The Elements of Statistical Learning: Data Mining, Inference, and Prediction*. Springer-Verlag, Berlin.
- HELFENSTEIN, F., E. DANCHIN, AND R. H. WAGNER. 2004. Assortative mating and sexual size dimorphism in Black-legged Kittiwakes. *Waterbirds* 27:350–354.
- HUBERTY, C. J. 1994. *Applied Discriminant Analysis*. Wiley, New York.
- IKO, N. M., S. J. DINSMORE, AND F. L. KNOFF. 2004. Evaluating the use of morphometric measurements from museum specimens for sex determination in Mountain Plovers (*Charadrius montanus*). *Western North American Naturalist* 64:492–496.
- ISAKSSON, A., M. WALLMAN, H. GÖRANSSON, AND M. G. GUSTAFSSON. 2008. Cross-validation and bootstrapping are unreliable in small sample classification. *Pattern Recognition Letters* 19:1960–1965.
- JAMES, F. C., AND C. E. MCCULLOCH. 1990. Multivariate analysis in ecology and systematics: Panacea or Pandora's box? *Annual Review of Ecology, Evolution, and Systematics* 21:129–166.
- JARRELL, M. G. 1994. A comparison of two procedures, the Mahalanobis distance and the Andrews-Pregibon statistic, for identifying multivariate outliers. *Research in the Schools* 1:49–58.
- JEFFREY, J. J., S. G. FANCY, G. D. LINDSEY, P. C. BANKO, T. K. PRATT, AND J. D. JACOBI. 1993. Sex and age identification of Palila. *Journal of Field Ornithology* 64:490–499.
- JOHNSON, R. A., AND D. W. WICHERN. 1992. *Applied Multivariate Statistical Analysis*, 3rd. ed. Prentice-Hall, Englewood Cliffs, New Jersey.
- KENWARD, B., C. RUTZ, A. A. S. WEIR, J. CHAPPELL AND A. KACELNIK. 2004. Morphology and sexual dimorphism of the New Caledonian Crow *Corvus moneduloides*, with notes on its behaviour and ecology. *Ibis* 146:652–660.
- LANCE, R. F., M. L. KENNEDY, AND P. L. LEBERG. 2000. Classification bias in discriminant function analyses used to evaluate putatively different taxa. *Journal of Mammalogy* 81:245–249.
- LOUGHEED, S. C., T. W. ARNOLD, AND R. C. BAILEY. 1991. Measurement error of external and skeletal variables in birds and its effect on principal components. *Auk* 108:432–436.
- MALLORY, M. L., AND M. R. FORBES. 2005. Sex discrimination and measurement bias in Northern Fulmars *Fulmarus glacialis* from the Canadian Arctic. *Ardea* 93:25–36.
- MANLY, B. F. J. 1994. *Multivariate Statistical Methods: A Primer*, 2nd ed. Chapman & Hall, London.
- MARDIA, K. V., J. T. KENT, AND J. M. BIBBY. 1979. *Multivariate Analysis*. Academic Press, New York.
- MARKS, S., AND O. J. DUNN. 1974. Discriminant functions when covariance matrices are unequal. *Journal of the American Statistical Association* 69:555–559.
- MCCARTHY, M. A. 2007. *Bayesian Methods for Ecology*. Cambridge University Press, Cambridge, United Kingdom.
- MCCRACKEN, K. G., D. C. PATON, AND A. D. AFTON. 2000. Sexual size dimorphism of the Musk Duck. *Wilson Bulletin* 112:457–466.
- MCGARIGAL, K., S. CUSHMAN, AND S. STAFFORD. 2000. *Multivariate Statistics for Wildlife and Ecology Research*. Springer-Verlag, New York.
- MILLER, W. J., AND F. H. WAGNER. 1955. Sexing mature columbiformes by cloacal characters. *Auk* 72:279–285.
- MØLLER, A. P., AND M. D. JENNIONS. 2001. Testing and adjusting for publication bias. *Trends in Ecology & Evolution* 16:580–586.
- MONCEAU, K. 2009. *Biologie des populations de tourterelles à queue carrée sur l'île de la Barbade: Apport de l'outil moléculaire*. Ph.D. dissertation, Université de Bourgogne, Dijon, France.
- MONCEAU, K., R. WATTIER, F.-X. DECHAUME-MONCHARMONT, S. MOTREUIL, AND F. CÉZILLY. 2011. Territoriality versus flocking in the Zenaida Dove (*Zenaida aurita*): Resource polymorphism revisited using morphological and genetic analyses. *Auk* 128:000–000.
- MORRISON, M. L. 1984. Influence of sample size on discriminant function analysis of habitat use by birds. *Journal of Field Ornithology* 55:330–335.

- MURPHY, M. T. 2007. A cautionary tale: Cryptic sexual size dimorphism in a socially monogamous passerine. *Auk* 124:515–525.
- NAKAGAWA, S., AND I. C. CUTHILL. 2007. Effect size, confidence interval and statistical significance: A practical guide for biologists. *Biological Reviews* 82:591–605.
- NORDHAUSEN, K., H. OJA, AND D. E. TYLER. 2008. Tools for exploring multivariate data: The package ICS. *Journal of Statistical Software* 28:1–31.
- OSBORNE, J. W., AND A. OVERBAY. 2004. The power of outliers (and why researchers should always check for them). *Practical Assessment, Research and Evaluation* 9(6).
- PALMER, A. R. 2000. Quasi-replication and the contract of error: Lessons from sex ratios, heritabilities and fluctuating asymmetry. *Annual Review of Ecology and Systematics* 31:441–480.
- PETRIDES, G. A. 1950. Notes on determination of sex and age in the Woodcock and Mourning Dove. *Auk* 67:357–360.
- PICARD, R. R., AND K. N. BERK. 1990. Data splitting. *American Statistician* 44:140–147.
- PIRAUX, F., AND R. PALM. 2001. Etude empirique des estimateurs des taux d'erreur en analyse discriminante. *Revue de statistique appliquée* 49:71–85.
- R DEVELOPMENT CORE TEAM. 2010. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna. [Online.] Available at cran.r-project.org/doc/manuals/refman.pdf.
- RASMUSSEN, J. L. 1988. Evaluating outlier identification tests: Mahalanobis D squared and Comrey D. *Multivariate Behavioral Research* 23:189–202.
- REMISIEWICZ, M., AND L. WENNERBERG. 2006. Differential migration strategies of the Wood Sandpiper (*Tringa glareola*)—Genetic analyses reveal sex differences in morphology and spring migration phenology. *Ornis Fennica* 83:1–10.
- ROBERTSON, B. C., AND N. J. GEMMELL. 2006. PCR-based sexing in conservation biology: Wrong answers from an accurate methodology? *Conservation Genetics* 7:267–271.
- RUBEGA, M. A. 1996. Sexual size dimorphism in Red-necked Phalaropes and functional significance of nonsexual bill structure variation for feeding performance. *Journal of Morphology* 228:45–60.
- RYDER, J. P. 1978. Sexing Ring-billed Gulls externally. *Bird-Banding* 49:218–222.
- SANTIAGO-ALARCON, D., AND P. G. PARKER. 2007. Sexual size dimorphism and morphological evidence supporting the recognition of two subspecies in the Galápagos Dove. *Condor* 109:132–141.
- SCARGLE, J. D. 2000. Publication bias: The “file-drawer” problem in scientific inference. *Journal of Scientific Exploration* 14:91–106.
- SCHROEDER, J., P. M. LOURENÇO, M. VAN DER VELDE, J. C. E. W. HOOIJMEIJER, C. BOTH, AND T. PIERSMA. 2008. Sexual dimorphism in plumage and size in Black-tailed Godwits *Limosa limosa limosa*. *Ardea* 96:25–37.
- SHEALER, D. A., AND C. M. CLEARY. 2007. Sex determination of adult Black Terns by DNA and morphometrics: Tests of sample size, temporal stability and geographic specificity in the classification accuracy of discriminant functions models. *Waterbirds* 30:180–188.
- SMITH, R. J. 1999. Statistics of sexual size dimorphism. *Journal of Human Evolution* 36:423–459.
- SNYDER, P., B. THOMPSON, AND P. SNYDER. 1991. Three reasons why stepwise regression methods should not be used by researchers. Pages 99–105 in *Advances in Educational Research: Substantive Findings, Methodological Developments* (B. Thompson, Ed.). JAI, Greenwich, Connecticut.
- STEVENS, J. 1992. *Applied Multivariate Statistics for the Social Sciences*, 2nd ed. Lawrence Erlbaum, Hillsdale, New Jersey.
- TABACHNICK, B. G., AND L. S. FIDELL. 2000. *Using Multivariate Statistics*, 4th ed. HarperCollins, New York.
- THOMPSON, B. 1995. Stepwise regression and stepwise discriminant analysis need not apply here: A guidelines editorial. *Educational and Psychological Measurement* 55:525–534.
- THOMPSON, B. 2001. Significance, effect sizes, stepwise methods, and other issues: Strong arguments move the field. *Journal of Experimental Education* 70:80–93.
- URA, T., N. AZUMA, S. HAYAMA, AND S. HIGASHI. 2005. Sexual dimorphism of Latham's Snipe (*Gallinago hardwickii*). *Emu* 105: 259–262.
- URFL, A. J., AND A. KALAM. 2006. Sexual size dimorphism and mating pattern in the Painted Stork (*Mycteria leucocephala*). *Waterbirds* 29:489–496.
- VENABLES, W. N., AND B. D. RIPLEY. 2002. *Modern Applied Statistics With S*, 4th ed. Springer-Verlag, Berlin.
- WEHBERG, S., AND M. SCHUMACHER. 2004. A comparison of nonparametric error rate estimation methods in classification problems. *Biometrical Journal* 46:35–47.
- WEIHS, C., U. LIGGES, K. LUEBKE, AND N. RAABE. 2005. klaR analyzing German business cycles. Pages 335–343 in *Data Analysis and Decision Support* (D. Baier, R. Decker, and L. Schmidt-Thieme, Eds.). Springer-Verlag, Berlin.
- WILLIAMS, B. K., AND K. TITUS. 1988. Assessment of sampling stability in ecological applications of discriminant analysis. *Ecology* 69:1275–1285.
- YEZERINAC, S. M., S. C. LOUGHEED, AND P. HANDFORD. 1992. Measurement error and morphometric studies: Statistical power and observer experience. *Systematic Biology* 41:471–482.
- ZAR, J. H. 1999. *Biostatistical Analysis*, 4th ed. Prentice Hall, Upper Saddle River, New Jersey.
- ZAVALAGA, C. B., AND R. PAREDES. 1997. Sex determination of adult Humboldt Penguins using morphometric characters. *Journal of Field Ornithology* 68:102–112.

Associate Editor: M. E. Hauber



Supplementary Online Material for

SEXING BIRDS USING DISCRIMINANT FUNCTION ANALYSIS: A CRITICAL APPRAISAL

FRANÇOIS-XAVIER DECHAUME-MONCHARMONT,^{1,3} KARINE MONCEAU,¹ AND FRANK CEZILLY^{1,2}

¹Université de Bourgogne, Equipe Ecologie Evolutive, UMR CNRS 5561 Biogéosciences, Dijon, France; and

²Institut Universitaire de France

TABLE S1. Summary of literature survey based on 141 articles. Discriminant function analysis (DFA) for sexing bird as function of family and species, age (A = adults, J = juveniles, AJ = adults and juveniles), sample size (number of birds used to calibrate the discriminant function), discriminant rate (proportion of birds correctly sexed), mean tarsus length, validation method (J = jack-knife, R = resubstitution, SS = sample splitting, NE = non-explained), variable reduction (A = automated stepwise, NA = non-automated variable reduction, N = no reduction, NE = non-explained), number of variables used for equation, authors, and year of publication.

Family/Species	Age	Sample size	Discriminant rate (%)	Tarsus length male (mm)	Tarsus length female (mm)	Validation method	Variable reduction	Number of variables used for equation	Authors	Year
Accipitridae										
<i>Aquila adalberti</i>	AJ	38	94.7	93.8	102.1	J	A	1	Ferrer and de le Court	1992
<i>Buteo jamaicensis</i>	J	121	97.0	88.2	85.2	NE	A	4	Donohue and Dufty	2006
<i>Buteo jamaicensis</i>	A	50	98.0	85.7	88.9	NE	A	2	Donohue and Dufty	2006
<i>Buteo jamaicensis</i>	A	49	100.0	10.8	12.5	SS	A	3	Pitzer et al.	2008
<i>Buteo jamaicensis</i>	J	51	90.3	10.6	12.1	SS	A	3	Pitzer et al.	2008
<i>Buteo jamaicensis</i>	J	139	90.0	10.7	12.1	SS	A	3	Pitzer et al.	2008
<i>Buteo lineatus</i>	J	25	100.0	7.8	8.5	SS	A	3	Pitzer et al.	2008
<i>Buteo swainsoni</i>	AJ	104	93.3	69.5	72.6	J	A	3	Sarasola and Negro	2004
<i>Haliaeetus albicilla</i>	J	182	96.4	12.8	14.8	R	A	1	Helander et al.	2007
<i>Hieraaetus fasciatus</i>	J	39	94.9	110.2	113.4	J	A	3	Palma et al.	2001
<i>Hieraaetus pennatus</i>	J	81	84.0	64.4	69.3	J	NA	4	Balbontin et al.	2001
<i>Hieraaetus pennatus</i>	A	41	100.0	64.1	69.4	J	NA	2	Balbontin et al.	2001
Alaudidae										
<i>Chersophilus duponti</i>	A	311	99.0	23.8	23.0	J	A	2	Vögeli et al.	2007
<i>Chersophilus duponti</i>	J	42	97.6	23.7	22.9	J	A	2	Vögeli et al.	2007
Alcedinidae										
<i>Todiramphus cinnamominus</i>	A	41	73.0	18.2	18.5	R	N	4	Kesler et al.	2006
Alcidae										
<i>Alca torda</i>	A	80	78.8	34.6	34.9	J	A	3	Grecian et al.	2003
<i>Alle alle</i>	A	141	70.2	20.3	20.2	J	A	1	Jakubas and Wojczulanis	2007
<i>Cerorhinca monocerata</i>	A	73	95.9	31.4	30.7	NE	A	2	Niizuma et al.	1999
<i>Fratercula cirrhata</i>	A	176	74.0	NA	NA	J	N	3	Williams et al.	2007
Ardeidae										
<i>Ardea alba</i>	A	76	89.5	153.2	145.2	J	A	2	Herring et al.	2008
Callaeidae										
<i>Callaeas cinerea</i>	A	130	82.6	68.6	64.5	NE	A	2	Flux and Innes	2001
Charadriidae										
<i>Charadrius montanus</i>	A	190	63.0	39.6	39.8	NE	A	1	Iko et al.	2004
Ciconiidae										
<i>Ciconia boyciana</i>	A	25	82.0	NA	NA	SS	N	2	Cheong et al.	2007

(continued)

³E-mail: fx.dechaume@u-bourgogne.fr

The following is supplementary to *The Auk*, Vol. 128, No. 1, pages 78–86.

The Auk, Vol. 128, Number 1, pages S1–S9. ISSN 0004-8038, electronic ISSN 1938-4254. © 2011 by The American Ornithologists' Union. All rights reserved. Please direct all requests for permission to photocopy or reproduce article content through the University of California Press's Rights and Permissions website, <http://www.ucpressjournals.com/reprintInfo.asp>. DOI: 10.1525/auk.2011.10129

TABLE S1. Continued.

Family/Species	Age	Sample size	Discriminant rate (%)	Tarsus length male (mm)	Tarsus length female (mm)	Validation method	Variable reduction	Number of variables used for equation	Authors	Year
Columbidae										
<i>Ducula goliath</i>	A	58	74.1	41.0	39.7	J	NA	6	Barré et al.	2003
<i>Zenaida galapagoensis</i>	A	105	97.5	NA	NA	J	A	3	Santiago-Alarcon and Parker	2007
Corvidae										
<i>Corvus brachyrhynchos</i>	A	74	91.9	58.9	56.2	NE	NA	3	Clark et al.	1991
<i>Corvus brachyrhynchos</i>	J	30	79.5	59.4	56.7	NE	NA	4	Clark et al.	1991
<i>Corvus frugilegus</i>	A	156	98.7	55.2	52.1	NE	NE	2	Green	1982
<i>Corvus monedula</i>	A	95	93.7	45.5	43.9	NE	NE	2	Green and Theobalt	1989
<i>Corvus moneduloides</i>	A	22	90.9	57.6	54.9	J	A	2	Kenward et al.	2004
<i>Cyanopica cyanus</i>	A	62	90.0	35.3	34.3	NE	A	3	Alarcos et al.	2007
<i>Cyanopica cyanus</i>	J	54	90.0	34.8	34.1	NE	A	2	Alarcos et al.	2007
<i>Pica pica</i>	A	71	88.7	50.4	48.0	NE	A	3	Reese and Kadlec	1982
<i>Pica pica</i>	AJ	243	93.8	NA	NA	NE	NE	5	Kavanagh	1988
<i>Pica pica</i>	J	105	86.5	50.3	48.2	NE	A	3	Lee et al.	2007
<i>Pica pica</i>	A	72	93.1	50.1	48.0	NE	A	3	Lee et al.	2007
<i>Pyrrhocorax pyrrhocorax</i>	A	51	100.0	55.7	51.6	SS	NE	2	Tella and Torre	1993
<i>Pyrrhocorax pyrrhocorax</i>	A	171	100.0	55.7	51.5	SS	N	2	Blanco et al.	1996
Emberizinae										
<i>Miliaria calandra</i>	A	103	96.1	NA	NA	R	N	2	Campos et al.	2005
<i>Passerculus sandwichensis</i>	J	119	94.1	21.2	20.7	NE	NE	2	Wheelwright et al.	1994
<i>Passerculus sandwichensis</i>	J	93	75.3	19.3	18.9	NE	NE	3	Wheelwright et al.	1994
Estrildidae										
<i>Lonchura striata</i>	A	25	84.0	13.4	12.4	NE	A	2	Mizuta et al.	2003
Falconidae										
<i>Falco peregrinus</i>	J	150	96.2	43.2	48.1	SS	NE	5	Hurley et al.	2007
Fringillidae										
<i>Hemignathus munroi</i>	A	30	93.0	25.1	24.0	J	A	2	Pratt et al.	1994
<i>Hemignathus munroi</i>	A	48	92.0	25.1	24.4	J	A	3	Pratt et al.	1994
<i>Pseudonestor xanthophrys</i>	A	51	97.0	23.2	21.4	J	A	1	Berlin et al.	2001
Furnariidae										
<i>Automolus ochrolaemus</i>	A	33	81.8	24.9	23.6	J	A	3	Winker et al.	1994
<i>Spartonoica maluioides</i>	A	35	77.0	19.6	19.4	J	A	4	Cardoni et al.	2009
<i>Xenops minutus</i>	A	31	90.3	14.2	14.1	J	A	1	Winker et al.	1994
<i>Xiphorhynchus flavigaster</i>	AJ	40	95.0	22.8	22.2	NE	A	2	Puebla-Olivares and Figueroa-Esquivel	2009
Haematopodidae										
<i>Haematopus bachmani</i>	A	119	88.2	51.6	53.1	J	A	4	Guzzetti et al.	2008
<i>Haematopus ostralegus</i>	A	1,195	90.7	NA	NA	R	N	4	Zwarts et al.	1996
<i>Haematopus ostralegus</i>	J	444	89.2	NA	NA	R	N	4	Zwarts et al.	1996
Hirundinidae										
<i>Hirundo rustica</i>	A	581	90.2	11.1	11.2	J	NE	3	Hermosell et al.	2007
<i>Hirundo rustica</i>	A	1,891	91.6	11.1	11.1	J	NE	3	Hermosell et al.	2007
Hydrobatidae										
<i>Oceanodroma furcata</i>	A	120	89.0	NA	NA	SS	NA	2	Boersma and Davies	1987
Laniidae										
<i>Lanius excubitor</i>	A	39	87.4	NA	NA	J	NA	2	Brady et al.	2009
Laridae										
<i>Anous stolidus</i>	A	49	90.0	NA	NA	R	A	2	Chardine and Morris	1989
<i>Chlidonias niger</i>	A	37	89.0	NA	NA	R	A	2	Stern and Jarvis	1991
<i>Chlidonias niger</i>	A	449	81.0	NA	NA	SS	A	2	Shealer and Cleary	2007
<i>Larus argentatus</i>	A	54	98.2	68.3	63.3	R	A	3	Shugart	1977
<i>Larus argentatus</i>	A	73	99.3	NA	NA	SS	A	2	Fox et al.	1981
<i>Larus argentatus</i>	A	480	96.8	NA	NA	NE	NA	6	Coulson et al.	1983
<i>Larus argentatus</i>	A	134	97.8	NA	NA	R	A	2	Migot	1986
<i>Larus argentatus</i>	A	316	92.0	71.1	66.0	SS	A	3	Evans et al.	1995
<i>Larus atricilla</i>	A	122	95.3	52.8	48.9	SS	A	2	Hanners and Patton	1985
<i>Larus atricilla</i>	A	76	93.0	50.5	47.3	SS	A	3	Evans et al.	1993
<i>Larus cachinnans</i>	A	181	100.0	71.0	65.1	J	A	4	Bosch	1996
<i>Larus californicus</i>	A	66	100.0	59.5	55.1	J	A	4	Schnell et al.	1985

(continued)

TABLE S1. Continued.

Family/Species	Age	Sample size	Discriminant rate (%)	Tarsus length male (mm)	Tarsus length female (mm)	Validation method	Variable reduction	Number of variables used for equation	Authors	Year
<i>Larus californicus</i>	A	491	99.2	57.6	53.3	SS	A	3	Rodriguez et al.	1996
<i>Larus californicus</i>	A	203	96.0	60.0	55.5	J	A	3	Herring et al.	2010
<i>Larus crassirostris</i>	A	237	96.6	57.6	53.2	NE	N	2	Chochi et al.	2002
<i>Larus delawarensis</i>	A	93	97.9	58.7	55.2	R	A	3	Shugart	1977
<i>Larus delawarensis</i>	A	59	95.0	66.2	61.4	SS	A	2	Ryder	1978
<i>Larus dominicanus</i>	A	100	97.0	NA	NA	SS	N	3	Torlaschi et al.	2000
<i>Larus fuscus</i>	A	121	97.5	NA	NA	NE	NA	6	Coulson et al.	1983
<i>Larus michahellis</i>	A	155	89.5	65.0	60.7	NE	A	4	Arizaga et al.	2008
<i>Larus michahellis</i>	A	67	97.0	74.2	68.9	J	A	3	Galarza et al.	2008
<i>Larus novaehollandiae</i>	A	165	90.1	45.6	43.8	NE	N	2	Mills	1971
<i>Larus ridibundus</i>	A	411	94.4	46.7	43.5	NE	A	2	Palomares et al.	1997
<i>Larus ridibundus</i>	J	143	90.2	46.6	44.0	NE	A	2	Palomares et al.	1997
<i>Onychoprion fuscatus</i>	A	63	77.8	25.5	24.8	NE	A	3	Reynolds et al.	2008
<i>Rissa tridactyla</i>	A	45	95.6	NA	NA	NE	NA	6	Coulson et al.	1983
<i>Rissa tridactyla</i>	A	303	84.2	37.0	35.4	SS	N	3	Jodice et al.	2000
<i>Rynchops niger</i>	A	50	100.0	33.5	28.9	J	A	1	Quinn	1990
<i>Rynchops niger</i>	A	78	97.9	33.9	29.8	NE	NA	2	Mariano-Jelicich et al.	2007
<i>Sterna caspia</i>	A	35	77.1	46.3	45.0	J	A	5	Quinn	1990
<i>Sterna caspia</i>	A	40	83.0	46.5	45.3	J	A	2	Ackerman et al.	2008
<i>Sterna forsteri</i>	A	84	86.0	24.7	24.3	J	A	2	Bluso et al.	2006
<i>Sterna hirundo</i>	A	105	80.0	24.0	24.1	NE	NA	4	Coulter	1986
<i>Sterna hirundo</i>	A	122	72.6	21.4	21.2	NE	A	3	Fletcher and Hamer	2003
<i>Sterna hirundo</i>	A	244	78.7	21.3	21.0	J	NA	2	Nisbet et al.	2007
<i>Sterna paradisaea</i>	A	71	71.6	16.8	16.6	NE	A	2	Fletcher and Hamer	2003
<i>Sterna paradisaea</i>	A	166	74.0	16.6	16.4	J	A	2	Devlin et al.	2004
Mimidae										
<i>Dumetella carolinensis</i>	A	242	78.0	NA	NA	SS	A	4	Suthers and Suthers	1990
<i>Mimodes graysoni</i>	A	35	100.0	37.8	35.9	J	NA	2	Martínez-Gómez and Curry	1998
Notiomystidae										
<i>Notiomystis cincta</i>	J	313	76.4	28.7	27.7	J	A	3	Thorogood et al.	2009
Otididae										
<i>Otis tarda</i>	J	165	98.2	125.5	108.2	NE	A	1	Martín et al.	2000
Paridae										
<i>Parus atricapillus</i>	A	314	93.7	16.9	16.3	SS	A	3	Desrochers	1990
Pelecanidae										
<i>Pelecanus erythrorhynchos</i>	A	188	97.0	120.4	111.0	J	A	2	Dorr et al.	2005
Phalacrocoracidae										
<i>Phalacrocorax albiventer</i>	A	84	96.5	65.0	62.2	NE	A	2	Malacalaza and Hall	1988
<i>Phalacrocorax atriceps</i>	A	188	94.0	69.0	65.1	J	A	2	Svagej and Quintana	2007
<i>Phalacrocorax auritus</i>	A	80	95.7	68.8	67.2	SS	A	3	Glahn and McCoy	1995
<i>Phalacrocorax bransfieldensis</i>	A	84	97.7	70.0	65.7	R	N	3	Casaux and Baroni	2000
<i>Phalacrocorax carbo</i>	A	51	96.1	NA	NA	R	A	3	Koffijberg and van Eerden	1995
<i>Phalacrocorax carbo</i>	A	81	95.1	69.4	65.1	J	NA	3	Liordos and Goutner	2008
<i>Phalacrocorax magellanicus</i>	A	84	86.0	54.0	52.8	NE	A	3	Quintana et al.	2003
Phoenicopteridae										
<i>Phoenicopus minor</i>	J	18	94.0	203.0	185.0	NE	A	2	Childress et al.	2005
<i>Phoenicopus minor</i>	A	40	93.0	238.0	210.0	NE	A	2	Childress et al.	2005
<i>Phoenicopus minor</i>	A	96	98.0	242.5	213.0	NE	A	2	Childress et al.	2005
Pipridae										
<i>Corapipo altera</i>	A	153	100.0	NA	NA	R	A	2	Mendenhall et al.	2010
<i>Manacus aurantiacus</i>	A	55	100.0	NA	NA	R	A	2	Mendenhall et al.	2010
Podicipedidae										
<i>Podiceps grisegena</i>	A	76	80.0	NA	NA	R	A	2	Kloskowski et al.	2006
<i>Podiceps nigricollis</i>	A	427	85.7	NA	NA	R	N	2	Jehl et al.	1998
Procellariidae										
<i>Calonectris diomedea</i>	J	94	93.7	NA	NA	J	N	4	Bretagnolle and Thibault	1995
<i>Calonectris diomedea</i>	A	211	95.0	54.2	52.8	NE	NA	2	Lo Valvo	2001
<i>Daption capense</i>	A	62	81.0	47.1	45.7	J	A	4	van Franeker and ter Braak	1993
<i>Fulmarus glacialis</i>	A	25	95.3	NA	NA	NE	N	2	Dunnet and Anderson	1961

(continued)

TABLE S1. Continued.

Family/Species	Age	Sample size	Discriminant rate (%)	Tarsus length male (mm)	Tarsus length female (mm)	Validation method	Variable reduction	Number of variables used for equation	Authors	Year
<i>Fulmarus glacialis</i>	A	247	98.0	55.8	51.7	J	A	4	van Franeker and ter Braak	1993
<i>Fulmarus glacialis</i>	A	32	97.0	56.0	52.5	J	A	4	van Franeker and ter Braak	1993
<i>Fulmarus glacialis</i>	A	63	95.0	54.1	49.8	J	A	2	Mallory and Forbes	2005
<i>Fulmarus glacialisoides</i>	A	130	89.0	54.3	51.9	J	A	4	van Franeker and ter Braak	1993
<i>Macronectes giganteus</i>	J	64	100.0	91.7	84.8	R	N	1	Copello et al.	2006
<i>Macronectes giganteus</i>	A	40	100.0	92.2	84.8	R	N	1	Copello et al.	2006
<i>Pagodroma nivea</i>	A	32	91.0	34.4	33.9	J	A	4	van Franeker and ter Braak	1993
<i>Pterodroma leucoptera</i>	A	209	66.2	30.4	30.2	J	N	3	O'Dwyer et al.	2006
<i>Pterodroma leucoptera</i>	J	206	64.6	30.6	30.3	J	N	2	O'Dwyer et al.	2006
<i>Puffinus carneipes</i>	A	102	90.0	56.1	54.9	J	A	3	Thalmann et al.	2007
<i>Puffinus mauretanicus</i>	A	52	90.0	50.2	49.0	J	NA	2	Genovart et al.	2003
<i>Puffinus yelkouan</i>	A	60	80.0	48.5	47.5	J	A	3	Bourgeois et al.	2007
<i>Thalassoica antarctica</i>	A	77	82.0	46.7	45.1	J	A	4	van Franeker and ter Braak	1993
<i>Thalassoica antarctica</i>	A	129	92.0	NA	NA	NE	NE	3	Lorentsen and Røv	1994
Psittacidae										
<i>Nestor notabilis</i>	A	61	85.5	42.6	41.5	NE	A	1	Bond et al.	1991
Rallidae										
<i>Fulica americana</i>	A	32	100.0	NA	NA	NE	NA	2	Boersma and Davies	1987
<i>Gallinula chloropus</i>	A	10	90.0	58.7	56.2	NE	N	7	Cucco et al.	1999
<i>Porphyrio mantelli</i>	A	37	89.2	97.3	90.7	J	A	2	Eason et al.	2001
<i>Rallus elegans</i>	A	26	100.0	61.0	54.2	SS	A	2	Perkins et al.	2009
<i>Rallus longirostris</i>	A	23	91.7	54.6	50.4	SS	A	3	Perkins et al.	2009
Scolopacidae										
<i>Calidris alba</i>	A	49	92.0	25.9	26.8	NE	A	2	Maron and Myers	1984
<i>Calidris alba</i>	A	42	86.0	NA	NA	NE	NA	2	Wood	1987
<i>Calidris alpina</i>	J	56	91.4	24.7	26.1	SS	NA	3	Meissner	2005
<i>Calidris alpina alpina</i>	A	80	98.8	24.7	26.1	J	NE	2	Meissner and Pilacka	2008
<i>Calidris alpina pacifica</i>	A	200	91.5	NA	NA	J	N	3	Brennan et al.	1984
<i>Calidris canutus</i>	A	112	75.9	31.6	32.5	R	N	6	Baker et al.	1999
<i>Calidris canutus</i>	A	90	80.0	NA	NA	R	N	4	Baker et al.	1999
<i>Calidris canutus</i>	A	85	67.1	NA	NA	R	N	4	Baker et al.	1999
<i>Calidris temminckii</i>	A	43	86.0	17.8	18.0	J	A	2	Lislevand et al.	2009
<i>Gallinago gallinago</i>	A	334	89.0	32.4	33.2	J	A	4	McCloskey and Thompson	2000
<i>Lymnocyrtus minimus</i>	A	299	99.0	24.5	24.4	J	A	4	Sikora and Dubiec	2007
<i>Phalaropus lobatus</i>	A	53	73.0	NA	NA	NE	N	4	Rubega	1996
Spheniscidae										
<i>Eudyptes chrysocome</i>	A	117	93.2	NA	NA	J	NA	2	Hull	1996
<i>Eudyptes schlegeli</i>	A	138	97.1	NA	NA	J	NA	2	Hull	1996
<i>Eudyptula minor</i>	A	43	94.8	NA	NA	SS	A	2	Renner and Davis	1999
<i>Eudyptula minor</i>	A	400	91.1	NA	NA	SS	A	1	Arnould et al.	2004
<i>Megadyptes antipodes</i>	A	60	93.0	NA	NA	SS	N	2	Setiawan et al.	2004
<i>Megadyptes antipodes</i>	J	50	86.0	NA	NA	SS	N	2	Setiawan et al.	2004
<i>Pygoscelis adeliae</i>	A	45	89.0	NA	NA	NE	A	3	Kerry et al.	1992
<i>Pygoscelis antarctica</i>	A	55	94.6	NA	NA	J	N	2	Amat et al.	1993
<i>Pygoscelis papua</i>	A	35	91.4	NA	NA	J	A	5	Renner et al.	1998
<i>Spheniscus humboldti</i>	A	223	97.0	NA	NA	SS	A	2	Zavalga and Paredes	1997
<i>Spheniscus magellanicus</i>	A	98	95.9	49.4	46.4	NE	A	3	Scolaro et al.	1983
<i>Spheniscus magellanicus</i>	A	37	92.0	NA	NA	SS	NA	1	Boersma and Davies	1987
<i>Spheniscus magellanicus</i>	J	266	78.0	NA	NA	J	NA	2	Bertelloti et al.	2002
<i>Spheniscus magellanicus</i>	A	331	97.0	NA	NA	J	NA	2	Bertelloti et al.	2002
Stercorariidae										
<i>Stercorarius parasiticus</i>	A	74	90.5	44.5	45.2	J	A	2	Phillips and Furness	1997
Strigidae										
<i>Aegolius funereus</i>	A	41	96.9	NA	NA	SS	N	4	Hayward and Hayward	1991
<i>Aegolius funereus</i>	AJ	135	70.0	NA	NA	NE	A	1	Hipkiss	2007
<i>Bubo bubo</i>	A	50	90.7	93.8	102.5	NE	N	4	del mar Delgado and Penteriani	2004
<i>Otus asio</i>	A	77	88.3	28.4	27.6	NE	A	3	Smith and Wiemeyer	1992

(continued)

TABLE S1. Continued.

Family/Species	Age	Sample size	Discriminant rate (%)	Tarsus length male (mm)	Tarsus length female (mm)	Validation method	Variable reduction	Number of variables used for equation	Authors	Year
<i>Strix aluco</i>	A	142	81.0	NA	NA	R	NE	2	Hardy et al.	1981
<i>Strix occidentalis</i>	A	133	90.2	60.2	61.6	J	NA	1	Blakesley et al.	1990
Sturnidae										
<i>Acridotheres javanicus</i>	A	69	72.5	37.7	37.1	J	A	5	Counsilman et al.	1994
<i>Acridotheres tristis</i>	A	90	84.0	38.3	36.7	J	A	7	Counsilman et al.	1994
<i>Sturnus roseus</i>	A	41	93.0	32.0	30.7	J	N	2	Zenatello and Kiss	2005
Tetraonidae										
<i>Lagopus leucurus</i>	A	49	87.0	NA	NA	SS	A	2	Gruys and Hannon	1993
<i>Lagopus leucurus</i>	J	49	89.8	NA	NA	SS	A	2	Gruys and Hannon	1993
Threskiornithidae										
<i>Eudocimus albus</i>	A	130	77.0	97.4	88.4	J	A	2	Herring et al.	2008
<i>Plegadis falcinellus</i>	A	198	84.9	95.4	85.2	J	A	2	Figuerola et al.	2006
Troglodytidae										
<i>Henicorhina prostheleuca</i>	AJ	83	95.2	22.4	21.3	J	A	4	Winker et al.	1996
<i>Thryothorus maculipictus</i>	AJ	62	95.2	21.9	20.8	J	A	3	Winker et al.	1996
<i>Troglodytes troglodytes</i>	A	85	96.0	NA	NA	NE	NA	2	Sweeney and Tatner	1996
Turdidae										
<i>Catharus bicknelli</i>	A	193	83.1	32.0	31.1	SS	A	2	Frey et al.	2008
<i>Catharus bicknelli</i>	J	128	79.1	32.0	31.1	SS	A	3	Frey et al.	2008
Tyrannidae										
<i>Empidonax virescens</i>	A	114	93.0	18.7	18.0	SS	A	2	Wilson	1999
Vireonidae										
<i>Hylophilus decurtatus</i>	A	44	97.7	16.9	16.7	NE	A	3	Winker et al.	1994
<i>Hylophilus ochraceiceps</i>	A	36	75.0	17.0	17.4	J	A	3	Winker et al.	1994

LITERATURE CITED

- ACKERMAN, J. T., J. Y. TAKEKAWA, J. D. BLUSO, J. L. YEE, AND C. A. EAGLES-SMITH. 2008. Gender identification of Caspian Terns using external morphology and discriminant function analysis. *Wilson Journal of Ornithology* 120:378–383.
- ALARCOS, S., C. DE LA CRUZ, E. SOLÍS, J. VALENCIA, AND M. J. GARCÍA-BAQUERO. 2007. Sex determination of Iberian Azure-winged Magpies *Cyanopica cyanus cooki* by discriminant analysis of external measurements. *Ring Migration* 23:211–216.
- AMAT, J. A., J. VIÑUELA, AND M. FERRER. 1993. Sexing Chinstrap Penguins (*Pygoscelis antarctica*) by morphological measurements. *Colonial Waterbirds* 16:213–215.
- ARIZAGA, J., A. ALDALUR, A. HERRERO, AND D. GALICIA. 2008. Sex differentiation of Yellow-legged Gull (*Larus michahellis lusitanicus*): The use of biometrics, bill morphometrics and wing tip coloration. *Waterbirds* 31:211–219.
- ARNOULD, J. P. Y., P. DANN, AND J. M. CULLEN. 2004. Determining the sex of Little Penguins (*Eudyptula minor*) in northern Bass Strait using morphometric measurements. *Emu* 104:261–265.
- BAKER, A. J., T. PIERSMA, AND A. D. GREENSLADE. 1999. Molecular vs. phenotypic sexing in Red Knots. *Condor* 101:887–893.
- BALBONTIN, J., M. FERRER, AND E. CASADO. 2001. Sex determination in Booted Eagles (*Hieraaetus pennatus*) using molecular procedures and discriminant function analysis. *Journal of Raptor Research* 35:20–23.
- BARRÉ, N., M. DE GARINE WICHATITSKY, R. LECOQ, AND J.-C. MAILLARD. 2003. Contribution to the knowledge of the New Caledonian Imperial Pigeon *Ducula goliath* (Gray 1859) with emphasis on sexual dimorphism. *Notornis* 50:155–160.
- BERLIN, K. E., J. C. SIMON, T. K. PRATT, P. E. BAKER, AND J. R. KOWALSKY. 2001. Age and sex determination of the Maui Parrotbill. *Journal of Field Ornithology* 72:12–21.
- BERTELLOTTI, M., J. L. TELLA, J. A. GODOY, G. BLANCO, M. G. FORERO, J. A. DONÁZAR, AND O. CEBALLOS. 2002. Determining sex of Magellanic Penguins using molecular procedures and discriminant functions. *Waterbirds* 25:479–484.
- BLAKESLEY, J. A., A. B. FRANKLIN, AND R. J. GUTIÉRREZ. 1990. Sexual dimorphism in Northern Spotted Owls from northwest California. *Journal of Field Ornithology* 61:320–327.
- BLANCO, G., J. L. TELLA, AND I. TORRE. 1996. Age and sex determination of monomorphic non-breeding Choughs: A long-term study. *Journal of Field Ornithology* 67:428–433.
- BLUSO, J. D., J. T. ACKERMAN, J. Y. TAKEKAWA, AND J. L. YEE. 2006. Sexing Forster's terns using morphometric measurements. *Waterbirds* 29:512–517.
- BOERSMA, P. D., AND E. M. DAVIES. 1987. Sexing monomorphic birds by vent measurements. *Auk* 104:779–783.
- BOND, A. B., K.-J. WILSON, AND J. DIAMOND. 1991. Sexual dimorphism in the Kea *Nestor notabilis*. *Emu* 91:12–19.
- BOSCH, M. 1996. Sexual size dimorphism and determination of sex in Yellow-legged Gulls. *Journal of Field Ornithology* 67:534–541.
- BOURGEOIS, K., C. CURÉ, J. LEGRAND, E. GÓMEZ-DÍAZ, E. VIDAL, T. AUBIN, AND N. MATHEVON. 2007. Morphological versus acoustic analysis: What is the most efficient method for sexing Yelkouan Shearwaters *Puffinus yelkouan*? *Journal of Ornithology* 148:261–269.

- BRADY, R. S., J. D. PARUK, AND A. J. KERN. 2009. Sexing adult Northern shrikes using DNA, morphometrics, and plumage. *Journal of Field Ornithology* 80:198–205.
- BRENNAN, L. A., J. B. BUCHANAN, C. T. SCHICK, S. G. HERMAN, AND T. M. JOHNSON. 1984. Sex determination of dunlins in winter plumage. *Journal of Field Ornithology* 55:343–348.
- BRETAGNOLLE, V., AND J.-C. THIBAUT. 1995. Method for sexing fledglings in Cory's Shearwaters and comments on sex-ratio variation. *Auk* 112:785–790.
- CAMPOS, F., M. HERNÁNDEZ, J. ARIZAGA, R. MIRANDA, AND A. AMEZCUA. 2005. Sex differentiation of Corn Buntings *Miliaria calandra* wintering in northern Spain. *Ring Migration* 22:159–162.
- CARDONI, D. A., J. E. MALDONADO, J. P. ISACCH, AND R. GREENBERG. 2009. Subtle sexual dimorphism in the Bay-capped Wren-spinetail (*Spartonotus maluroides*; Furnariidae) uncovered through molecular sex determination. *Ornitología Neotropical* 20:347–355.
- CASAUX, R., AND A. BARONI. 2000. Sexual size dimorphism in the Antarctic shag. *Waterbirds* 23:489–493.
- CHARDINE, J. W., AND R. D. MORRIS. 1989. Sexual size dimorphism and assortative mating in the Brown Noddy. *Condor* 91:868–874.
- CHEONG, S., H.-C. SUNG, AND S.-R. PARK. 2007. A new method for sexing Oriental White Storks. *Journal of Field Ornithology* 78:329–333.
- CHILDRESS, B., D. HARPER, B. HUGHES, AND C. FERRIS. 2005. Sex determination in the Lesser Flamingo (*Phoenicopterus minor*) using morphological measurements. *Ostrich* 76:148–153.
- CHOCHI, M., Y. NIZUMA, AND M. TAKAGI. 2002. Sexual differences in the external measurements of Black-tailed Gulls breeding on Rishiri Island, Japan. *Ornithological Sciences* 1:163–166.
- CLARK, R. G., P. C. JAMES, AND J. B. MORARI. 1991. Sexing adult and yearling American Crows by external measurements and discriminant analysis. *Journal of Field Ornithology* 62:132–138.
- COPELLO, S., F. QUINTANA, AND G. SOMOZA. 2006. Sex determination and sexual size-dimorphism in Southern Giant Petrels (*Macronectes giganteus*) from Patagonia, Argentina. *Emu* 106:141–146.
- COULSON, J. C., C. S. THOMAS, J. E. L. BUTTERFIELD, N. DUNCAN, P. MONAGHAN, AND C. SHEDDEN. 1983. The use of head and bill length to sex live gulls Laridae. *Ibis* 125:549–557.
- COULTER, M. C. 1986. Assortative mating and sexual dimorphism in the Common Tern. *Wilson Bulletin* 98:93–100.
- COUNSILMAN, J. J., K. NEE, A. K. JALIL, AND W. L. KENG. 1994. Discriminant analysis of morphometric characters as a means of sexing Mynas. *Journal of Field Ornithology* 65:1–7.
- CUCCO, M., G. LINGUA, D. BOCCHIO, C. ACQUARONE, AND G. MALACARNE. 1999. Sex identification in the Moorhen (*Gallinula chloropus*) by flow cytometry and morphometric analysis. *Italian Journal of Zoology* 66:1–6.
- DEL MAR DELGADO, M., AND V. PENTERIANI. 2004. Gender determination of Eurasian Eagle-owls (*Bubo bubo*) by morphology. *Journal of Raptor Research* 38:375–377.
- DESROCHERS, A. 1990. Sex determination of Black-capped chickadees with discriminant analysis. *Journal of Field Ornithology* 61:79–84.
- DEVLIN, C. M., A. W. DIAMOND, AND G. W. SAUNDERS. 2004. Sexing Arctic Terns in the field and laboratory. *Waterbirds* 27:314–320.
- DONOHUE, K. C., AND A. M. DUFTY, JR. 2006. Sex determination of Red-tailed Hawks (*Buteo jamaicensis calurus*) using DNA analysis and morphometrics. *Journal of Field Ornithology* 77:74–79.
- DORR, B., D. T. KING, J. B. HARREL, P. GERARD, AND M. G. SPALDING. 2005. The use of culmen length to determine sex of the American White Pelican. *Waterbirds* 28:102–106.
- DUNNET, G. M., AND A. ANDERSON. 1961. A method for sexing living Fulmars in the hand. *Bird Study* 8:119–126.
- EASON, D., C. D. MILLAR, A. CREE, J. L. HALVERSON, AND D. M. LAMBERT. 2001. A comparison of five methods for assignment of sex in the Takahe (*Aves: Porphyrio mantelli*). *Journal of Zoology* 253:281–292.
- EVANS, D. R., P. M. CAVANAGH, T. W. FRENCH, AND B. G. BLODGET. 1995. Identifying the sex of Massachusetts Herring Gulls by linear measurements. *Journal of Field Ornithology* 66:128–132.
- EVANS, D. R., E. M. HOOPES, AND C. R. GRIFFIN. 1993. Discriminating the sex of Laughing Gulls by linear measurements. *Journal of Field Ornithology* 64:472–476.
- FERRER, M., AND C. DE LE COURT. 1992. Sex Identification in the Spanish Imperial Eagle. *Journal of Field Ornithology* 63:359–364.
- FIGUEROLA, J., L. GARCIA, A. J. GREEN, F. IBAÑEZ, M. MAÑEZ, J. L. DEL VALLE, H. GARRIDO, J. L. ARROYO, AND R. RODRÍGUEZ. 2006. Sex determination in Glossy Ibis chicks based on morphological characters. *Ardeola* 53:229–235.
- FLETCHER, K. L., AND K. C. HAMER. 2003. Sexing terns using biometrics: The advantage of within-pair comparisons. *Bird Study* 50:78–83.
- FLUX, I., AND J. INNES. 2001. A field technique for determining the sex of North Island Kokapo (*Callaeas cinerea wilsoni*). *Notornis* 48:217–223.
- FOX, G. A., C. R. COOPER, AND J. P. RYDER. 1981. Predicting the sex of Herring Gulls by using external measurements. *Journal of Field Ornithology* 52:1–9.
- VAN FRANEKER, J. A., AND J. F. TER BRAAK. 1993. A generalized discriminant for sexing Fulmarine Petrels from external measurements. *Auk* 110:492–502.
- FREY, S. J. K., C. C. RIMMER, K. P. MCFARLAND, AND S. MENU. 2008. Identification and sex determination of Bicknell's Thrushes using morphometric data. *Journal of Field Ornithology* 79:408–420.
- GALARZA, A., J. HIDALGO, G. OCIO, AND P. RODRIGUEZ. 2008. Sexual size dimorphism and determination of sex in Atlantic Yellow-legged Gulls *Larus michahellis lusitanicus* from northern Spain. *Ardeola* 55:41–47.
- GENOVART, M., M. MCMINN, AND D. BOWLER. 2003. A discriminant function for predicting sex in the Balearic Shearwater. *Waterbirds* 26:72–76.
- GLAHN, J. F., AND R. B. MCCOY. 1995. Measurements of wintering Double-crested Cormorants and discriminant models of sex. *Journal of Field Ornithology* 66:299–304.
- GRECIAN, V. D., A. W. DIAMOND, AND J. W. CHARDINE. 2003. Sexing Razorbills *Alca torda* breeding at Machias Seal Island, New Brunswick, Canada, using discriminant function analysis. *Atlantic Seabirds* 5:73–80.
- GREEN, P. T. 1982. Sexing Rooks *Corvus frugilegus* by discriminant analysis. *Ibis* 124:320–324.
- GREEN, P. T., AND C. M. THEOBALD. 1989. Sexing birds by discriminant analysis: Further considerations. *Ibis* 131:442–447.

- GRUYS, R. C., AND S. J. HANNON. 1993. Sex determination of hunter-killed and depredated Willow Ptarmigan using a discriminant analysis. *Journal of Field Ornithology* 64:11–17.
- GUZZETI, B. M., S. L. TALBOT, D. F. TESSLER, V. A. GILL, AND E. C. MURPHY. 2008. Secrets in the eyes of Black Oystercatchers: A new sexing technique. *Journal of Field Ornithology* 79:215–223.
- HANNERS, L. A., AND S. R. PATTON. 1985. Sexing Laughing Gulls using external measurements and discriminant analysis. *Journal of Field Ornithology* 56:158–164.
- HARDY, A. R., G. J. M. HIRONS, P. I. STANLEY, AND L. W. HUSON. 1981. Sexual dimorphism in size of Tawny Owls (*Strix aluco*): A method for sexing in field studies. *Ardea* 69:181–184.
- HAYWARD, G. D., AND P. H. HAYWARD. 1991. Body measurements of Boreal Owls in Idaho and a discriminant model to determine sex of live specimens. *Wilson Bulletin* 103:497–500.
- HELANDER, B., F. HAILER, AND C. VILÀ. 2007. Morphological and genetic sex identification of White-tailed Eagle *Haliaeetus albicilla* nestlings. *Journal of Ornithology* 148:435–442.
- HERMOSELL, I. G., J. BALBONTIN, A. MARZAL, M. REVIRIEGO, AND F. DE LOPE. 2007. Sex determination in Barn Swallows *Hirundo rustica* by means of discriminant analysis in two European populations. *Ardeola* 54:93–100.
- HERRING, G., D. E. GAWLIK, AND J. M. BEERENS. 2008. Sex determination for the Great Egret and White Ibis. *Waterbirds* 31:298–303.
- HERRING, G., J. T. ACKERMAN, C. A. EAGLES-SMITH, AND J. Y. TAKEKAWA. 2010. Sexing California Gulls using morphometrics and discriminant function analysis. *Waterbirds* 33:79–85.
- HIPKISS, T. 2007. Can migrating Tengmalm's Owls *Aegolius funereus* be reliably sexed in autumn using simple morphometric measurements? *Ring Migration* 23:201–204.
- HULL, C. L. 1996. Morphometric indices for sexing adult Royal *Eudyptes schlegeli* and Rockhopper *E. chrysocome* penguins at Macquarie Island. *Marine Ornithology* 24:23–27.
- HURLEY, V. G., F. HOGAN, J. G. WHITE, AND R. COOKE. 2007. A morphological model for sexing nestling Peregrine Falcons (*Falco peregrinus macropus*) verified through genetic analysis. *Wildlife Research* 34:54–58.
- IKO, N. M., S. J. DINSMORE, AND F. L. KNOPF. 2004. Evaluating the use of morphometric measurements from museum specimens for sex determination in Mountain Plovers (*Charadrius montanus*). *Western North American Naturalist* 64:492–496.
- JAKUBAS, D., AND K. WOJCZULANIS. 2007. Predicting the sex of Dovekies by discriminant analysis. *Waterbirds* 30:92–96.
- JEHL, J. R., A. E. HENRY, AND S. I. BOND. 1998. Sexing Eared Grebes by bill measurements. *Colon. Waterbird* 21:98–100.
- JODICE, P. G. R., R. B. LANCTOT, V. A. GILL, D. D. ROBY, AND S. A. HATCH. 2000. Sexing adult Black-legged Kittiwakes by DNA, behavior, and morphology. *Waterbirds* 23:405–415.
- KAVANAGH, B. 1988. Discriminating the sex of Magpies *Pica pica* from morphological data. *Ring Migration* 9:83–90.
- KENWARD, B., C. RUTZ, A. A. S. WEIR, J. CHAPPELL, AND A. KACELNIK. 2004. Morphology and sexual dimorphism of the New Caledonian Crow *Corvus moneduloides*, with notes on its behaviour and ecology. *Ibis* 146:652–660.
- KERRY, K. R., D. J. AGNEW, J. R. CLARKE, AND G. D. ELSE. 1992. Use of morphometric parameters for the determination of sex of Adélie Penguins. *Wildlife Research* 19:657–664.
- KESLER, D. C., I. F. LOPES, AND S. M. HAIG. 2006. Sex determination of Pohnpei Micronesian Kingfishers using morphological and molecular genetic techniques. *Journal of Field Ornithology* 77:229–232.
- KLOSKOWSKI, J., P. GRELA, J. KROGULEC, M. GASKA, AND M. T. CHORZEWSKI. 2006. Sexing Red-necked Grebes *Podiceps grisegena* by molecular techniques and morphology. *Acta Ornithologica* 41:176–180.
- KOFFIJBERG, K., AND M. R. VAN EERDEN. 1995. Sexual dimorphism in the cormorant *Phalacrocorax carbo sinensis*: Possible implications for differences in structural size. *Ardea* 83:37–46.
- LEE, S.-I., H.-J. JANG, S.-H. EO, AND J.-C. CHOE. 2007. Sexual size dimorphism and morphological sex determination in the Black-billed Magpie in South Korea (*Pica pica sericea*). *Journal of Ecology and Field Biology* 30:195–199.
- LIORDOS, V., AND V. GOUTNER. 2008. Sex determination of Great Cormorants (*Phalacrocorax carbo sinensis*) using morphometric measurements. *Waterbirds* 31:203–210.
- LISLEVAND, T., G. MARTHINSEN, AND J. T. LIFJELD. 2009. Sex differences in body size and body condition in breeding Temminck's Stints *Calidris temminckii*. *Journal of Ornithology* 150:299–302.
- LO VALVO, M. 2001. Sexing adult Cory's Shearwater by discriminant analysis of body measurements on Linosa Island (sicilian channel), Italy. *Waterbirds* 24:169–174.
- LORENTSEN, S.-H., AND N. RØV. 1994. Sex determination of Antarctic Petrels *Thalassoica antarctica* by discriminant analysis of morphometric characters. *Polar Biology* 14:143–145.
- MALACALAZA, V. E., AND M. A. HALL. 1988. Sexing adult King Cormorants (*Phalacrocorax albiventer*) by discriminant analysis. *Colonial Waterbird* 11:32–37.
- MALLORY, M. L., AND M. R. FORBES. 2005. Sex discrimination and measurement bias in Northern Fulmars *Fulmarus glacialis* from the Canadian Arctic. *Ardea* 93:25–36.
- MARIANO-JELICICH, R., E. MADRID, AND M. FAVEROL. 2007. Sexual dimorphism and diet segregation in the Black Skimmer *Rynchops niger*. *Ardea* 95:115–124.
- MARON, J. L., AND J. P. MYERS. 1984. A description and evaluation of two techniques for sexing wintering Sanderlings. *Journal of Field Ornithology* 55:336–342.
- MARTÍN, C. A., J. C. ALONSO, J. A. ALONSO, M. B. MORALES, AND C. PITRA. 2000. An approach to sexing young Great Bustards *Otis tarda* using discriminating analysis and molecular techniques. *Bird Study* 47:147–153.
- MARTINEZ-GÓMEZ, J. E., AND R. L. CURRY. 1998. Distinguishing sex of Socorro Mockingbirds by body measurements. *Ornitología Neotropical* 9:103–110.
- MCCLOSKEY, J. T., AND J. E. THOMPSON. 2000. Aging and sexing Common Snipe using discriminant analysis. *Journal of Wildlife Management* 64:960–969.
- MEISSNER, W. 2005. Sex determination of juvenile Dunlins migrating through the Polish Baltic region. *Journal of Field Ornithology* 76:368–372.
- MEISSNER, W., AND L. PILACKA. 2008. Sex identification of adult Dunlins *Calidris alpina alpina* migrating in autumn through Baltic region. *Ornis Fennica* 85:135–139.
- MENDENHALL, C. D., C. H. SEKERCIOGLU, AND F. O. BRENES. 2010. Using interpubic distance for sexing Manakins. *Journal of Field Ornithology* 81:49–63.

- MIGOT, P. 1986. Le goéland argenté *Larus argentatus argenteus* Brehm en Bretagne: Caractéristiques biométriques des reproducteurs. *Alauda* 54:269–278.
- MILLS, J. A. 1971. Sexing Red-billed Gulls from standard measurements. *New Zealand Journal of Marine Freshwater Research* 5:326–328.
- MIZUTA, T., H. YAMADA, R.-S. LIN, Y. YODOGAWA, AND K. OKANOYA. 2003. Sexing Whiterumped Munias in Taiwan, using morphology, DNA and distance calls. *Ornithological Science* 2:97–102.
- NIIZUMA, Y., A. TAKAHASHI, M. KUROKI, AND Y. WATANUKI. 1999. Sexing by external measurements of adult Rhinoceros Auklets breeding on Teuri Island. *Japanese Journal of Ornithology* 48:145–150.
- NISBET, I. C. T., E. S. BRIDGE, P. SZCZYS, AND B. J. HEIDINGER. 2007. Sexual dimorphism, female-female pairs, and test for assortative mating in Common terns. *Waterbirds* 30:169–179.
- O'DWYER, T. W., D. PRIDDEL, N. CARLILE, J. A. BARTLE, AND W. A. BUTTEMER. 2006. An evaluation of three field techniques for sexing Gould's Petrels (*Pterodroma leucoptera*) (Procellariidae). *Emu* 106:245–252.
- PALMA, L., S. MIRA, P. CARDIA, P. BEJA, T. GUILLEMAUD, N. FERRAND, M. L. CANCELA, AND L. C. DA FONSECA. 2001. Sexing Bonelli's eagle nestlings: Morphometrics versus molecular techniques. *Journal of Raptor Research* 35:187–193.
- PALOMARES, L. E., B. E. ARROYO, J. MARCHAMALO, J. J. SAINZ, AND B. VOSLAMBER. 1997. Sex- and age-related biometric variation of Black-headed Gulls *Larus ridibundus* in Western European populations. *Bird Study* 44:310–317.
- PERKINS, M., S. L. KING, S. TRAVIS, AND J. LINScombe. 2009. Use of morphometric measurements to differentiate between species and sex of King and Clapper Rails. *Waterbirds* 32:579–584.
- PHILLIPS, R. A., AND R. W. FURNESS. 1997. Predicting the sex of Parasitic Jaegers by discriminant analysis. *Colonial Waterbird* 20:14–23.
- PITZER, S., J. HULL, H. B. ERNEST, AND A. C. HULL. 2008. Sex determination of three raptor species using morphology and molecular techniques. *Journal of Field Ornithology* 79:71–79.
- PRATT, T. K., S. G. FANCY, C. K. HARADA, G. D. LINDSEY, AND J. D. JACOBI. 1994. Identifying sex and age of Akiapolaau. *Wilson Bulletin* 106:421–430.
- PUEBLA-OLIVAREZ, F., AND E. M. FIGUEROLA-ESQUIVEL. 2009. Sexual dimorphism in Ivory-billed Woodcreepers (*Xiphorhynchus flavigaster*) in Mexico. *Journal of Ornithology* 150:755–760.
- QUINN, J. S. 1990. Sexual size dimorphism and parental care patterns in a monomorphic and a dimorphic larid. *Auk* 107:260–274.
- QUINTANA, F., G. SOMOZA, M. UHART, C. CASSARÁ, P. GANDINI, AND E. FRERE. 2003. Sex determination of adult Rock Shags by molecular sexing and morphometric parameters. *Journal of Field Ornithology* 74:370–375.
- REESE, K. P., AND J. A. KADLEC. 1982. Determining the sex of Black-billed Magpies by external measurements. *Journal of Field Ornithology* 53:417–418.
- RENNER, M., AND L. S. DAVIS. 1999. Sexing Little Penguins *Eudyptula minor* from Cook Strait, New Zealand using discriminant function analysis. *Emu* 99:74–79.
- RENNER, M., J. VALENCIA, L. S. DAVIS, D. SAEZ, AND O. CIFUENTES. 1998. Sexing adult Gentoo Penguins in Antarctica using morphometrics. *Colonial Waterbird* 21:444–449.
- REYNOLDS, S. J., G. R. MARTIN, L. L. WALLACE, C. P. WEARN, AND B. J. HUGHES. 2008. Sexing Sooty Terns on Ascension Island from morphometric measurements. *Journal of Zoology* 274:2–8.
- RODRIGUEZ, E. F., B. H. PUGESSEK, AND K. L. DIEM. 1996. A sexing technique for California Gulls breeding at Bamforth Lake, Wyoming. *Journal of Field Ornithology* 67:519–524.
- RUBEGA, M. A. 1996. Sexual size dimorphism in Red-necked Phalaropes and functional significance of non sexual bill structure variation for feeding performance. *Journal of Morphology* 228:45–60.
- RYDER, J. P. 1978. Sexing Ring-billed Gulls externally. *Bird Banding* 49:218–222.
- SANTIAGO-ALARCON, D., AND P. G. PARKER. 2007. Sexual size dimorphism and morphological evidence supporting the recognition of two subspecies in the Galápagos Dove. *Condor* 109:132–141.
- SARASOLA, J. H., AND J. J. NEGRO. 2004. Gender determination in the Swainson's Hawk (*Buteo swainsoni*) using molecular procedures and discriminant function analysis. *Journal of Raptor Research* 38:357–361.
- SCHNELL, G. D., G. L. WORTHEN, AND M. E. DOUGLAS. 1985. Morphometric assessment of sexual dimorphism in skeletal elements of California Gulls. *Condor* 87:484–493.
- SCOLARO, J. A., M. A. HALL, AND I. M. XIMÉNEZ. 1983. The Magellanic Penguin (*Spheniscus magellanicus*): Sexing adults by discriminant analysis of morphometric characters. *Auk* 100:221–224.
- SETIAWAN, A. N., J. T. DARBY, AND D. M. LAMBERT. 2004. The use of morphometric measurements to sex Yellow-eyed Penguins. *Waterbirds* 27:96–101.
- SHEALER, D. A., AND C. M. CLEARY. 2007. Sex determination of adult Black Terns by DNA and morphometrics: Tests of sample size, temporal stability and geographic specificity in the classification accuracy of discriminant functions models. *Waterbirds* 30:180–188.
- SHUGART, G. W. 1977. A method for externally sexing Gulls. *Bird Banding* 48:118–121.
- SIKORA, A., AND A. DUBIEC. 2007. Sex identification of Jack Snipe *Limnocryptes minimus* by discriminant analysis of morphometric measurements. *Ardea* 95:125–133.
- SMITH, D. G., AND S. N. WIEMEYER. 1992. Determining sex of Eastern Screech-owls using discriminant function analysis. *Journal of Raptor Research* 26:24–26.
- STERN, M. A., AND R. L. JARVIS. 1991. Sexual dimorphism and assortative mating in Black Terns. *Wilson Bulletin* 103:266–271.
- SUTHERS, H. B., AND D. D. SUTHERS. 1990. Aging and sexing Gray Catbirds by external characteristics. *North American Bird Bander* 15:45–52.
- SVAGELJ, W. S., AND F. QUINTANA. 2007. Sexual size dimorphism and sex determination by morphometric measurements in breeding Imperial Shags (*Phalacrocorax atriceps*). *Waterbirds* 30:97–102.
- SWEENEY, J. J., AND P. TATNER. 1996. Sexing Wrens *Troglodytes troglodytes indigenus* using morphological measurements and discriminant analysis. *Bird Study* 43:342–350.
- TELLA, J. L., AND I. TORRE. 1993. Sexual size dimorphism and determination of sex Chough (*Pyrrhocorax pyrrhocorax*). *Journal of Ornithology* 134:187–190.
- THALMANN, S., B. BAKER, M. HINDELL, M. C. DOUBLE, AND R. GALES. 2007. Using biometric measurements to determine gender of Flesh-footed Shearwaters, and their application as a

- tool in long-line by-catch management and ecological field studies. *Emu* 107:231–238.
- THOROGOOD, R., D. BRUNTON, AND I. CASTRO. 2009. Simple techniques for sexing nestling Hihi (*Notiomystis cincta*). *New Zealand Journal of Zoology* 36:115–121.
- TORLASCHI, C., P. GANDINI, E. FRERE, AND R. M. PECK. 2000. Predicting the sex of Kelp Gulls by external measurements. *Waterbirds* 23:518–520.
- VÖGELI, M., D. SERRANO, J. L. TELLA, M. MENDEZ, AND J. A. GODOY. 2007. Sex determination Dupont's Lark *Chersophilus duponti* using molecular sexing and discriminant functions. *Ardeola* 54:69–79.
- WHEELWRIGHT, N. T., G. TRUSSELL, J. P. DEVINE, AND R. ANDERSON. 1994. Sexual dimorphism and population sex ratios in juvenile Savannah Sparrows. *Journal of Field Ornithology* 65:520–529.
- WILLIAMS, C. T., S. D. KILDAW, AND C. L. BUCK. 2007. Sex-specific differences in body condition indices and seasonal mass loss in Tufted Puffins. *Journal of Field Ornithology* 78:369–378.
- WILSON, R. R. 1999. Sex determination of the Acadian Flycatcher using discriminant analysis. *Journal of Field Ornithology* 70:514–519.
- WINKER, K., J. T. KLIKA, AND G. VOELKER. 1996. Sexual size dimorphism in birds from southern Veracruz, Mexico. 2. *Thryothorus maculipectus* and *Henicorhina [Leucosticta] prostheleuca*. *Journal of Field Ornithology* 67:236–251.
- WINKER, K., G. A. VOELKER, AND J. T. KLIKA. 1994. A morphometric examination of sexual dimorphism in the *Hylophilus*, *Xenops*, and an *Automolus* from southern Veracruz, Mexico. *Journal of Field Ornithology* 65:307–323.
- WOOD, A. G. 1987. Discriminating the sex of Sanderling *Calidris alba*: Some results and their implications. *Bird Study* 34:200–204.
- ZAVALAGA, C. B., AND R. PAREDES. 1997. Sex determination in Humboldt Penguins using morphometric characters. *Journal of Field Ornithology* 68:102–112.
- ZENATELLO, M., AND J. B. KISS. 2005. Biometrics and sex identification of the Rose-coloured Starling *Sturnus roseus*. *Ring Migration* 22:163–166.
- ZWARTS, L., J. B. HULSCHER, K. KOOPMAN, AND P. M. ZEGERS. 1996. Discriminating the sex of Oystercatchers *Haematopus ostralegus*. *Ardea* 84A:1–12.